

# Specimen Label

2,4-D CHOLINE SALT

GROUP

4

HERBICIDE



## Enlist One®

with COLEX-D® technology

### HERBICIDE

TM®Trademarks of Corteva Agriscience and its affiliated companies

**This labeling expires on January 11, 2029.**

**Do not use or distribute this product after January 11, 2029.**

**For control of emerged annual and perennial broadleaf weeds, use as a preplant, preemergence and postemergence herbicide on Enlist® corn, soybeans and cotton. Enlist herbicides with Colex-D® technology are the ONLY 2,4-D containing products authorized and specifically labeled for use with Enlist crops.**

**Use as a non-selective burndown; chemical fallow; use as a preplant and preemergence herbicide on non-Enlist corn, and use as a preplant herbicide on non-Enlist soybeans.**

**Do not allow contact of herbicide with foliage of desirable plants and trees because severe injury or destruction may result.**

**Approved for use only in certain geographical areas. Read and follow all label instructions.**

**Active Ingredient(s):**

2,4-Dichlorophenoxyacetic acid,  
choline salt ..... 55.7%

**Other Ingredients.....** 44.3%

**Total .....** 100.0%

2,4-dichlorophenoxyacetic acid equivalent – 38% – 3.8 lb/gal

### FIRST AID

If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling poison control center (1-800-222-1222) or doctor, or going for treatment. You may also contact 1-800-992-5994, for emergency medical treatment information.	

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

EPA Reg. No. 62719-695

#### Keep Out of Reach of Children

#### WARNING

May be fatal if swallowed. Causes substantial but temporary eye injury. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Do not get in eyes or on clothing. Avoid contact with skin.

#### Personal Protective Equipment (PPE)

All mixers, loaders, applicators, flaggers, and handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks, plus
- Waterproof gloves
- Protective eyewear (goggles, faceshield, or safety glasses)
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate

See engineering controls for additional requirements.

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift or runoff may adversely affect aquatic invertebrates, sensitive wetland environments, and non-target plants. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift and runoff.

This product is moderately toxic to bees on an acute basis, and may cause chronic risk to pollinators or other terrestrial invertebrates. Do not apply this product to blooming vegetation or if bees or other pollinating insects are visiting the treatment area.

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several days to weeks after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of 2,4-D from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

2,4-D is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. Application around a cistern or well may result in contamination of drinking water or groundwater.

#### Physical and Chemical Hazards

Spray solutions of this product must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic lined containers.

**Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel containers or spray tanks.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

#### Endangered Species

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered

species and certain threatened species, under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email [ESPP@epa.gov](mailto:ESPP@epa.gov). You must use the Bulletin valid for the month in which you will apply the product.

**Report ecological incidents:** To report ecological incidents, including mortality, injury, or harm to non-target plants and animals, call 1-855-ENLIST-1 (1-855-365-4781).

#### **Tank-Mixing Instructions:**

Enlist One may only be tank-mixed with products that have been tested and found not to adversely affect the spray drift properties of Enlist One. A list of those products may be found at [Enlist.com/TankMix](http://Enlist.com/TankMix).

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### **DO NOT TANK-MIX ANY PRODUCT WITH Enlist One unless:**

1. You check the list of tested products found not to adversely affect the spray drift properties of Enlist One at [Enlist.com/TankMix](http://Enlist.com/TankMix) no more than 7 days before applying Enlist One; and
2. The product you tank-mix with Enlist One is identified on that list of tested products.

### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, faceshield, or safety glasses)

### **Product Information**

Enlist One herbicide is a systemic herbicide that is intended for control of emerged annual and perennial broadleaf weeds. Enlist One is designed to be applied to crops containing Enlist™ traits. These are patented genes that provide tolerance to Enlist One. Certain other uses (e.g. use as a non-selective burndown; chemical fallow; use as a preplant herbicide on non-Enlist soybeans, and a preplant or preemergence herbicide on non-Enlist corn) are also permitted as specified on this label. Corn, soybeans, and cotton without the Enlist trait will be seriously damaged by foliar applications of Enlist One.

When this product is applied as directed and under the circumstances described, it controls annual and perennial broadleaf weeds listed in this label.

**Time to Symptoms on Susceptible Plants:** Initial symptoms include drooping leaves and epinasty, which typically occurs within 24 hours of foliar treatment. This is followed by chlorosis, necrosis, further leaf/stem malformation and growth inhibition. Complete death and desiccation of susceptible plants occurs within 3-5 weeks.

**Stage of Broadleaf Weeds:** Apply when weeds are less than 6 inches in height. Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the annual and perennial tables for specific weeds. When treating weeds with disease or insect damage, weeds heavily covered with dust, or weeds under poor growing conditions, reduced weed control may result.

**Rainfastness:** Enlist One is rainfast within 4 hours following application. See Use Restrictions for application prior to rainfall and/or irrigation.

**Spray Coverage:** For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

**Mode of Action:** 2,4-D, the active ingredient in this product, mimics the naturally occurring plant auxins and overloads the plant's auxin balance affecting vital processes, such as cell division and elongation, resulting in abnormal growth and plant death.

**Limited Soil Activity:** Though some suppression of annual weeds emerging soon after application may occur, optimum control is achieved when the majority of weeds are emerged at the time of application.

**Biological Degradation:** Degradation of this product is primarily a biological process carried out by soil microbes.

### **Herbicide Resistance Management**

2,4-D, the active ingredient in this product, is a Group 4 herbicide (synthetic auxin). Some naturally occurring weed biotypes that are tolerant (resistant) to 2,4-D may exist due to genetic variability in a weed population. Where resistant biotypes exist, the repeated use of herbicides with the same modes of action can lead to the selection for resistant weeds. Certain agronomic practices delay or reduce the likelihood that resistant weed populations will develop and can be utilized to manage weed resistance once it occurs.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued availability of this product depends on the successful management of the weed resistance program; therefore, it is very important to perform the following actions.

To aid in the prevention of developing weeds resistant to this product, the following steps must be followed:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Apply full rates of Enlist One in combination with another herbicide with a different mode of action and overlapping spectrum (See Tank Mix Instructions). Choose the rate for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in weed species.
- Report any incidence of non-performance of this product against a particular weed species to your Corteva retailer, representative or call 1-855-ENLIST-1(1-855-365-4781).
- Suspected Resistance: Indicators of suspected herbicide resistance include (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of uncontrolled plants of a particular weed species; and (3) surviving plants mixed with controlled individuals of the same species. Likely resistant weeds are assumed to be present if any of these criteria are met.
- If resistance is suspected, treat weed escapes with an herbicide having a mode of action other than Group 4 and/or use non-chemical methods to remove escapes, as practicable, with the goal of preventing further seed production.

Additionally, users should follow as many of the following herbicide resistance management practices as practicable:

- Use a broad spectrum soil-applied herbicide with other modes of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative modes of action.
- Rotate the use of this product with non-Group 4 herbicides.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Thoroughly clean plant and soil residues from equipment before leaving fields suspected to contain resistant weeds.
- Avoid using more than two in-crop applications of Enlist One and any other Group 4 herbicide within a single growing season unless in conjunction with another mode of action herbicide with overlapping spectrum.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

Contact the local agricultural extension service, Corteva representative, ag retailer or crop consultant for further guidance on weed control practices as needed.

### **Susceptible Plants**

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Do not allow contact of herbicide with foliage, green stems, exposed non-woody roots of crops, desirable plants; including trees and cotton without the Enlist trait,

because severe injury or destruction may result. Small amounts of spray drift that may not be visible may injure susceptible broadleaf plants.

**Before making an application, please refer to your state's sensitive crop registry (if available) to identify any commercial specialty or certified organic crops that may be located nearby.**

**At the time of application, the wind cannot be blowing toward adjacent commercially grown tomatoes and other fruiting vegetables (EPA crop group 8), cucurbits (EPA crop group 9, including pumpkins, melons, and cucumbers), grapes, tobacco and cotton.**

## Spray Drift Management

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product. The applicator is responsible for avoiding off-site drift. Be aware of nearby non-target sites and environmental conditions.

### Do not aerially apply this product.

#### Nozzle Selection

The listing of nozzles and pressures on Enlist.com/Nozzles specifies which nozzles are allowed for use when applying Enlist One herbicide. Do not use any nozzle and pressure combination not specifically allowed by the listing on Enlist.com/Nozzles.

#### Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

#### Groundboom Application

Use the minimum boom height based upon the nozzle manufacturer's directions. Do not exceed 24 inches in height above the canopy. Spray drift potential increases as boom height increases. Spray drift can be minimized if nozzle height is not greater than the maximum height specified by the nozzle manufacturer for the nozzle selected.

#### Wind Speed

Do not apply at wind speeds greater than 15 mph. Wind speeds can vary during application. For best results apply when wind speeds are between 3 and 10 mph.

#### Temperature and Humidity


When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

#### Temperature Inversions

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground

source generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

#### Protection of Sensitive Areas



Applicator must maintain a 30-foot downwind buffer (in the direction in which the wind is blowing) from any area except:

1. Roads, paved or gravel surfaces.
2. Planted agricultural fields. (Except those crops listed in the "Susceptible Plants" section.)
3. Agricultural fields that have been prepared for planting.
4. Areas covered by the footprint of a building, shade house, silo, feed crib, or other man-made structure with walls and/or a roof.

To maintain the required downwind buffer zone:

- Measure wind direction prior to the start of any swath that is within 30 feet of a sensitive area.
- No application swath can be initiated in, or into an area that is within 30 feet of a sensitive area if the wind direction is towards the sensitive area.

#### State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

#### Management of Runoff

A variety of factors including soil type, slope, and weather conditions (e.g., rainfall) can influence volume and intensity of water running off the treated field. The applicator must evaluate all factors and make appropriate adjustments when applying this product. Land management, field condition and application practices that reduce, to the maximum extent practicable, runoff from treated fields, must be implemented by land managers/users of this product.

To reduce the potential for runoff and avoid off field impact from treated fields to maximum extent practicable, applicator must plan/schedule applications to maximize time between an application of this product and anticipated rainfall (or planned irrigation). Application must take place no less than 48 hours prior to irrigation or predicted rainfall (by NOAA/National Weather Service, or other similar forecasting service).

For land with **Hydrologic Soil Groups\* A & B**: The land manager/applicator must effectively implement measures in the following tables to equal a **minimum of 4 credits**.

For land with **Hydrologic Soil Groups\* C & D**: The land manager/applicator must effectively implement the measures in the following tables to equal a **minimum of 6 credits**.

Mitigation Measures	Credits
<b>Reduce number of applications</b> - Reduced number of applications of Enlist products per year. Applications may be made at any time during crop development but must maintain a minimum 12-day retreatment interval.	3 applications 0 2 applications 2 1 application 4
<b>Residue Tillage Management:</b> no-till, strip-till, ridge-till, and mulch-till	4
<b>Vegetative Filter Strips</b>	30 ft off-field vegetative buffer on down slope HSG A or B 2 HSG C or D 0 100 ft off-field vegetative buffer on down slope HSG A or B 4 HSG C or D 1
<b>Field border:</b> border with dense vegetative stands with a minimum width of 30 ft.	2
<b>Cover Crop</b>	2
<b>Vegetative Barrier:</b> Permanent strips of dense vegetation along the contours of the field with a minimum width of 3 ft.	2
<b>Contour Buffer Strips or Terrace</b>	2
<b>Grassed Waterway</b>	2
<b>Water and Sediment Basin</b>	1
<b>Contour Farming or Contour Strip Cropping</b>	1

\*Hydrologic Soil Group (HSG) definitions: A = Sand, loamy sand, or sandy loam; B = Sandy clay loam; C = Silt loam or loam; D = Clay loam, silty clay loam, sandy clay, silty clay or clay.

Applicators/Land Managers must meet minimum criteria described for each mitigation measure as outlined on Enlist.com/mitigationmeasures to receive credits.

## Sprayer Clean-Out

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Completely drain the spray system, including pump, lines and spray boom, for at least 5 minutes.
2. Fill the spray tank with clean water to at least 10% of the total tank volume and circulate the solution through the entire system so that all internal surfaces are contacted for at least 15 minutes to complete the first rinse of the application equipment. Spray the solution out of the spray tank through the boom.
3. Completely drain the spray system, including lines and spray boom, for at least 5 minutes; remove and clean filters and strainers.
4. During the second rinse, fill the container with clean water to at least 10% of the total tank volume. The addition of tank cleaning agents may be used at the manufacturer's specified rates. Circulate the solution through the entire system for at least 15 to 20 minutes. Let the solution stand for several hours, preferably overnight. Spray the solution out of the spray tank through the boom.
5. Completely drain the spray system, including lines and spray boom, for at least 5 minutes.
6. Fill the container with clean water to at least 10% of the total tank volume and circulate the solution through the entire system so that all internal surfaces are contacted for at least 15 minutes to complete the third rinse of the application equipment. Spray the solution out of the spray tank through the boom.
7. Completely drain the spray system, remove nozzle tips and strainers and clean them separately.

## Tank Mix Sequence Procedures

Find the list of qualified tank mix partners at [Enlist.com/TankMix](http://Enlist.com/TankMix).

- Start with a clean sprayer before mixing a load with Enlist One herbicide.
- Recommended water carrier volume with Enlist One is 10-15 gallons per acre.
- Do not use less than 10 gallons.
- Consult [Enlist.com/Nozzles](http://Enlist.com/Nozzles) or Enlist Product Use Guide for qualified nozzles and corresponding pressure ranges.
- For more tips on sprayer set up, visit [Enlist.com](http://Enlist.com).

## Mixing Steps

Begin with half-tank full of water carrier. Begin agitation and continue throughout mixing process. Add products in order, one at a time allowing time for thorough mixing before adding the next product:

1. AMS / water conditioning agents.
2. Pre-slurry water-soluble packets.
3. Wettable powders/dry flowables.
4. Compatibility agents
5. Liquid flowables.
6. Capsule suspension (CS) or suspension emulsion (SE).
7. Emulsifiable concentrate (EC).
8. Soluble liquids (SL)
9. Crop Oil Concentrate (COC), NIS, or other adjuvants
10. Top off with water carrier.

## Application Equipment and Application Methods

**Chemigation:** Do not apply this product through any type of irrigation system.

**Aerial Application:** Do not aerially apply this product.

Apply Enlist One with the following application equipment: Apply spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes.

## Ground Broadcast Spray

Boom, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment. Use the minimum boom height based upon the nozzle manufacturer's specifications. Spray drift potential is increased as boom height increases. Spray drift can be minimized if nozzle height is not greater than maximum height recommended by nozzle manufacturer for the nozzle selected. Do not apply greater than 24" above the crop canopy. Find the listing of nozzles and pressures on [Enlist.com/Nozzles](http://Enlist.com/Nozzles). This website specifies which nozzles are allowed for use when applying Enlist herbicides.

Use the specified rates of this product as a broadcast spray. As the density of weeds increases, increase spray volume within the specified range to ensure complete coverage. Check for even distribution of spray droplets.

## USES

Applications may be made to control any weeds listed in the annual and perennial tables.

### Precautions:

- The use directions are based upon a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence.
- In no-till and stale seedbed systems, a preplant burndown application is recommended to control existing weeds prior to crop emergence.
- Prior to application, follow all directions in Spray Drift Management and Management of Runoff sections for all applications.

### Restrictions

- **Crop Rotation/Plant-back Interval:** For any crop not listed in this section, do not apply less than 30 days prior to planting.
- Do not apply this product when soil is saturated or at field capacity, or when a storm event likely to produce runoff from the treated area is forecasted (by NOAA/National Weather Service, or other similar forecasting service) to occur within 48 hours following application.
- Do not irrigate treated fields within 48 hours of application.
- Do not apply more than 6 pints of Enlist One per acre per year (a combined total of 3 lbs 2,4-D a.e. per acre per year).
- Do not apply less than 12 days between applications.
- Enlist One is approved for use in the following states: Alabama, Arizona, Arkansas, Colorado, Delaware, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Virginia, West Virginia and Wisconsin. Do not use in any other state.
- All users must check *Bulletins Live!* Two prior to application as described in Endangered Species Advisory/Protection Requirements.
- **Endangered Species Advisory/Protection Requirements:** This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures controlling the product use relevant to your location for the protection of Endangered Species. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email [ESPP@epa.gov](mailto:ESPP@epa.gov). You must use the Bulletin valid for the month in which you will apply the product.
- **Do not use** Enlist One in the following counties:

State	County Restrictions
Florida	Brevard, Broward, Charlotte, Collier, DeSoto, Glades, Hardee, Hendry, Highlands, Hillsborough, Indian River, Lee, Manatee, Martin, Miami-Dade, Okeechobee, Orange, Osceola, Palm Beach, Polk, Sarasota, and St. Lucie

## Enlist Corn

These directions are for use on corn containing the Enlist trait. Information on crop varieties containing these traits may be obtained from your seed supplier.

### Carriers and Spray Volumes

Apply in a broadcast spray volume of water ranging from 10 to 15 gallons per acre for best results. Do not apply less than 10 gallons total spray volume per acre. Do not substitute water with nitrogen or other fertilizer solutions as carrier.

### Preplant (Burndown) Through Preemergence

Make a single application of 1.5 to 2.0 pints of Enlist One per acre. Use the upper end of the rate range for less susceptible weeds, more mature weeds, or weeds under stress. Refer to Weed Control section for specific weed height and use rate information. Apply any time before or after planting, but before corn emerges, to control weed seedlings or existing cover crops.

### Postemergence

Apply 2.0 pints of Enlist One per acre. Apply when weeds are no larger than 6 inches and corn is no larger than V8 growth stage or 30 inches (free standing) tall, whichever occurs first. For corn heights 30 to 48 inches (free standing), apply only using ground application equipment using drop nozzles aligned to avoid spraying into the whorl of corn plants. Make one to two applications with a minimum of 12 days between applications.

### Precautions:

- Application may result in temporary, cosmetic injury in the form of spotting or temporary plant leaning. This crop response will not affect long-term crop development or yield.
- Prior to application, follow all directions in Spray Drift Management and Management of Runoff sections for all applications.

**Restrictions:**

- **Preharvest Interval:** Do not apply within 30 days of forage harvest.
- Do not apply more than one preemergence application and no more than two postemergence applications per year. Using fewer applications will result in credits to satisfy the mitigation requirements (See Management of Runoff section).
- Do not apply more than 3 lbs 2,4-D a.e. per acre per year.
- Do not apply more than 2.0 pints (1 lb 2,4-D a.e.) of Enlist One per acre per application.
- Do not apply more than 6.0 pints (3 lbs a.e. 2,4-D) of Enlist One per acre per year.
- Do not apply Enlist One as a preharvest application or as an application to corn later than the V8 stage of corn that is more than 48 inches (free standing).
- Do not aerially apply this product.

**Corn – Not Containing the Enlist Trait**

**Labeled Crops:** Field corn, seed corn, sweet corn, popcorn

**Carriers and Spray Volumes**

Apply in a broadcast spray volume of water ranging from 10 to 15 gallons per acre for best results. Do not apply less than 10 gallons total spray volume per acre. Do not substitute water with nitrogen or other fertilizer solutions as carrier.

**Preplant (Burndown) Through Preemergence**

Make a single application of 1.5 to 2.0 pints of Enlist One per acre. Use the upper end of the rate range for less susceptible weeds, more mature weeds, or weeds under stress. Refer to Weed Control section for specific weed height and use rate information. Apply any time before or after planting, but before corn emerges, to control weed seedlings or existing cover crops.

**Postharvest**

Allow weeds to regrow after any damage incurred during harvest and recover from environmental stress before applying this product. Apply 2 pints of Enlist One per acre. Apply prior to heading of grass weeds and, if possible, before broadleaf weeds are more than 24 inches tall.

**Precautions:**

- For best results, do not apply to light sandy soils as a preplant or preemergence application.
- Using fewer applications will result in credits to satisfy the mitigation requirements (see Management of Runoff section).
- Prior to application, follow all directions in Spray Drift Management and Management of Runoff sections for all applications.

**Restrictions:**

- Do not aerially apply this product.
- Do not apply more than 2.0 pints (1 lb 2,4-D a.e.) of Enlist One per acre per application.
- Do not apply more than 4.0 pints (2 lbs 2,4-D a.e.) of Enlist One per acre per year.

**Enlist Soybeans**

These directions are for use with soybean containing the Enlist trait.

Information on crop varieties containing these traits may be obtained from your seed supplier.

**Carriers and Spray Volumes**

Apply in a broadcast spray volume of water ranging from 10 to 15 gallons per acre for best results. Do not apply less than 10 gallons total spray volume per acre. Do not substitute water with nitrogen or other fertilizer solutions as carrier.

**Preplant (Burndown) Through Preemergence**

Make a single application of 1.5 to 2.0 pints of Enlist One per acre. Use the upper end of the rate range for less susceptible weeds, more mature weeds, or weeds under stress. Refer to Weed Control section for specific weed height and use rate information. Apply any time before or after planting, but before soybean emerges, to control weed seedlings or existing cover crops.

**Postemergence**

Apply 2.0 pints of Enlist One per acre. Apply when weeds are no larger than 6 inches and any time after soybean emergence through the R1 growth stage. Make one to two applications with a minimum of 12 days between applications.

**Precaution:**

- Prior to application, follow all directions in Spray Drift Management and Management of Runoff sections for all applications.

**Restrictions:**

- **Preharvest Interval:** Do not apply within 50 days of harvest.
- Do not graze treated soybean.
- Do not harvest for forage or hay.
- Do not apply more than one preemergence application and no more than two postemergence applications per year. Using fewer

applications will result in credits to satisfy the mitigation requirements (See Management of Runoff section).

- Do not apply after R1 growth stage.
- Do not apply more than 3 lbs 2,4-D a.e. per acre per year.
- Do not apply more than 2.0 pints (1 lb a.e. 2,4-D) of Enlist One per acre per application.
- Do not apply more than 6.0 pints (3 lbs a.e. 2,4-D) of Enlist One per acre per year.
- Do not aerially apply this product.

**Soybean - Not Containing the Enlist Trait****Carriers and Spray Volumes**

Apply in a broadcast spray volume of water ranging from 10 to 15 gallons per acre for best results. Do not apply less than 10 gallons total spray volume per acre. Do not substitute water with nitrogen or other fertilizer solutions as carrier

**Preplant (Burndown)**

Apply up to 1.0 pints of Enlist One per acre no less than 7 days or up to 2.0 pints per acre, not less than 14 days prior to planting soybeans. See Precautions and Restrictions in this section. Refer to Weed Control section for specific weed height and use rate information.

**Precautions:**

- **Note:** Unacceptable injury to soybeans planted in treated fields may occur. Whether soybean injury occurs and the extent of such injury depends upon weather (temperature and rainfall) from herbicide application until soybean emergence, and agronomic factors, such as the amount of weed vegetation and previous crop residue present at the time of application. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.
- Do not disturb treated soil through tillage between application and planting of soybeans.
- Do not apply Enlist One as a preplant application in soybeans unless soybean injury is acceptable, including possible stand loss and/or yield reductions.
- Prior to application, follow all directions in Spray Drift Management and Management of Runoff sections for all applications.

**Restrictions:**

- Do not disturb treated soil through tillage between application and planting of soybeans.
- Do not use on sandy soils with less than 1% organic matter.
- In treated fields, plant soybean seed as deep as practicable, but not less than 1 inch deep. Adjust the planter, if necessary, to ensure that planted seed is adequately covered.
- Do not make more than one application per year regardless of the amount of product applied.
- During the growing season following application, do not replant treated fields with crops other than those labeled for use with 2,4-D.
- Do not apply more than 2.0 pints (1 lb a. e. 2,4-D) of Enlist One per acre per year.
- Do not aerially apply this product.

**Enlist Cotton**

These directions are for use on Cotton containing the Enlist trait.

Information on crop varieties containing these traits may be obtained from your seed supplier.

**Carriers and Spray Volumes**

Apply in a broadcast spray volume of water ranging from 10 to 15 gallons per acre for best results. Do not apply less than 10 gallons total spray volume per acre. Do not substitute water with nitrogen or other fertilizer solutions as carrier.

**Preplant (Burndown) Through Preemergence**

Make a single application of 1.5 to 2.0 pints of Enlist One per acre. Use the upper end of the rate range for less susceptible weeds, more mature weeds, or weeds under stress. Refer to Weed Control section for specific weed height and use rate information. Apply any time after planting, but before cotton emerges, to control weed seedlings or existing cover crops.

**Postemergence**

Apply 2.0 pints of Enlist One per acre. Apply when weeds are no larger than 6 inches and any time after cotton emergence up to first white bloom. Refer to Weed Control section for specific weed height and use rate information. Make one to two postemergence applications with a minimum of 12 days between applications.

**Precaution:**

- Prior to application, follow all directions in Spray Drift Management and Management of Runoff sections for all applications.

**Restrictions:**

- **Preharvest Interval:** Do not apply within 30 days of harvest.
- Do not graze treated cotton.
- Do not harvest for forage or hay.

- Do not apply more than one preemergence application and no more than two postemergence applications per year. Using fewer applications will result in credits to satisfy the mitigation requirements (See Management of Runoff section).
- Do not apply after first white bloom.
- Do not apply more than 2 pints Enlist One per acre per application (1 lb a.e. 2,4-D per acre).
- Do not apply more than 6 pints of Enlist One per acre per year (3 lbs a.e. 2,4-D per acre per year).
- Do not apply more than 3 lbs 2,4-D a.e. per acre per year
- Do not aerially apply this product.

## Fallow Systems to be Planted to Corn, Soybeans or Cotton

### Fallow

This product may be applied during the fallow period prior to planting or emergence of any crop listed on this label. This product may be used as a substitute for tillage to control annual weeds in fallow fields. Broadcast treatments will control or suppress many perennial weeds in fallow fields. Apply 2 pints of Enlist One per acre. Do not apply more than 6 pints of Enlist One per acre within the calendar year. Refer to Weed Control section for specific weed height and use rate information. Plant only labeled crops within 30 days following application.

### Carriers and Spray Volumes

Apply in a broadcast spray volume of water ranging from 10 to 15 gallons per acre for best results. Do not apply less than 10 gallons total spray volume per acre. Do not substitute water with nitrogen or other fertilizer solutions as carrier.

### Precaution:

- Prior to application, follow all directions in Spray Drift Management and Management of Runoff sections for all applications.

### Restrictions

- Do not aerially apply this product.
- Do not apply more than 2 pints per acre per application (1 lb a.e. 2,4-D per acre).
- Do not apply more than 6 pints per acre per year (a combined total of 3 lbs 2,4-D a.e. per acre per year).
- Do not apply more than 3 lbs 2,4-D a.e. per acre per year.

### Weed Control

Apply this product to actively growing weeds. Apply 2.0 pints of this product per acre for all postemergence uses with Enlist crops. Apply when weeds are no larger than 6 inches. Water carrier volumes of 10 to 15 gallons per acre are recommended for best results. Do not apply less than 10 gallons total spray volume per acre. Best control will be achieved when this product is applied in combination with another broad spectrum herbicide having a different mode of action (see Tank Mix Instructions).

Hard to control weeds, such as Palmer amaranth, may require a total program approach including soil applied residual herbicide(s) followed by a single or sequential post herbicide application.

Below-ground portions of perennial weeds may not be completely controlled with single applications and follow-up applications may be required if regrowth occurs.

## Controlled Weeds Table:

### Annual Weeds:

anoda, spurred	falseflax, smallseed	nightshade, black
bittercress	fiddleneck, field	nightshade, hairy
bitterweed	pennycress	Palmer amaranth <sup>1</sup>
broomweed, common	filareefleabane, annual	pepperweed
burdock	fleabane, hairy ( <i>Conyza bonariensis</i> ) <sup>1</sup>	pigweed, redroot
buttercup	fleabane, rough <sup>1</sup>	pigweed, smooth
carpetweed	geranium, Carolina	prickly lettuce
cinquefoil, common	groundcherry, hemp	puncturevine
cinquefoil, rough	sesbania	purslane
cocklebur	horseweed/	pusley, Florida
copperleaf,	maretail ( <i>Conyza canadensis</i> ) <sup>1</sup>	radish, wild
hophornbeam	jewelweed	ragweed, common
copperleaf, Virginia	jimsonweed	ragweed, giant
croton, Texas	lambsquarters	Russian thistle
croton, woolly	London rocket	salsify, common
dayflower, Benghal	mallow, Venice	salsify, western
devilsclaw (unicorn plant)	morningglory ( <i>Pomoea</i> spp.)	shepherd's-purse
dwarfandelion	mustard, tansy	sicklepod
eclipta	mustard, tumble	smartweed,
eveningprimrose,	mustard, wild	ladysthumb
common		smartweed,
falsedandelion		Pennsylvania
		sowthistle, annual

## Controlled Weeds Table: (Cont.)

### Annual Weeds:

Spanishneedles	thistle, bull	vetch
sunflower	thistle, musk	waterhemp <sup>1</sup>
sweetclover	velvetleaf	
teaweed/prickly sida	vervain	

<sup>1</sup>Hard to control weeds, such as Palmer amaranth or waterhemp, may require a total program approach including soil-applied residual herbicide(s) followed by a single or sequential post herbicide application.

### Perennial Weeds:

alfalfa	dock	pokeweed, common
artichoke, Jerusalem	dogbane	pennywort
aster, many flowered	garlic, wild	plantains
bindweed, field	hawkweed, orange	ragwort, tansy
bindweed, hedge	healall	sowthistle, perennial
blueweed, Texas	ironweed	thistle, Canada
catnip	ivy, ground	waterplantain
chicory	loco, bigbend	wormwood
cress, hoary	nettles	
dandelion	onion, wild	

## Storage and Disposal

Do not contaminate water, food, feed or seed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place. Store in original container. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### Nonrefillable containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

### Refillable containers larger than 5 gallons:

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

### Nonrefillable containers 5 gallons or larger:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and

## Storage and Disposal (Cont.)

continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

## Revisions:

1. Adjust titles of Base label and booklet cover.
2. Moved First Aid higher in both base label and booklet.
3. Added header "User Safety Requirements".
4. On base label added Header "Directions for Use" and added statement "It is a violation to Federal Law to..." In booklet moved this statement up in label.
5. Moved Storage and Disposal section to above the Warranty Section.
6. Removed statement "No till will result...of Runoff Section)." from 2nd bullet of overall Precautions.
7. Added to Uses Restrictions the following two statements:
  - 8. "Do not harvest or feed...unless otherwise specified.
  - 9. "Crop Rotation/Plant-back Interval:.....prior to planting."
10. Overall Use Restrictions added, "For Sub-County restrictions in Texas, Alabama, Georgia and Tennessee refer to" the restriction that started with "Endangered Species Advisory/Protection Requirements.."
11. Remove County restrictions.
12. Add timing restriction of no application between October 1 and May 15 for Wilson county in Tennessee.
13. To Soybean – Not Containing the Enlist Trait added the restriction, "Do not disturb treated soil...planning of soybeans."
14. Added statement, "All users must check Bulletins Live! Two prior to application as described in Endangered Species Advisory/Protection Requirements."
15. Across all Enlist Crops (corn, soybean, cotton) revised the restriction regarding number of applications to read as follows: Do not apply more than one preemergence application and no more than two postemergence applications per year. Using fewer applications will result in credits to satisfy the mitigation requirements (See Management of Runoff section).
16. Changed season to year within restrictions soybean not with Enlist trait.

## Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

## Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

## Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. Corteva Agriscience will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by Corteva Agriscience. To the extent permitted by law, all such risks associated with non-directed use shall be assumed by buyer and/or user.

## Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Corteva Agriscience be liable for consequential, incidental or special damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™®Trademarks of Corteva Agriscience and its affiliated companies

## Produced for

**Corteva Agriscience LLC**

**9330 Zionsville Road**

**Indianapolis, IN 46268**

Label Code: CD02-427-022

Replaced Label: CD02-427-021

EPA accepted 01/25/2024