

# TIGRIS™ PS CLETHODIM

## HERBICIDE

**ACTIVE INGREDIENT:**

\*Clethodim ..... 26.4%

**OTHER INGREDIENTS:** ..... 73.6%

**TOTAL:** ..... 100.0%

Contains Petroleum Distillates

\*(E)-2-[[(2-chloro-2-propenyl)-oxy]imino]propyl-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one

Contains 2.0 lbs. clethodim per gallon.

### KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

#### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Immediately call a poison control center or doctor. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. **DO NOT** give any liquid to the person. **DO NOT** give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**HOTLINE NUMBERS:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378, Mon.-Fri. 8:00 a.m. to 12:00 p.m. Pacific Time or your poison control center at 1-800-222-1222.

**NOTE TO PHYSICIAN:** Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid, which can cause pneumonitis. If ingested, probable mucosal damage may contraindicate the use of gastric lavage.

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

EPA Reg. No.: 92647-31

EPA Est. No.: 70989-M0-001 (OS)

05905-IA-001 (HD)

Letter(s) in the lot number correspond  
to letter(s) following the EPA Est. No.

Net Contents: 2.5 Gallons


**Manufactured For:**

Tigris, LLC  
P.O. Box 250  
10025 Hwy. 264 Alternate  
Middleses, NC 27557



6 44216 04945 5

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### WARNING

Causes substantial but temporary eye irritation. Avoid contact with skin. **DO NOT** get in eyes, on skin, or on clothing. Harmful if swallowed or inhaled. Avoid breathing vapors or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves including Barrier Laminote or Viton  $\geq$  14 mils
- Shoes plus socks
- Protective eyewear

#### USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

##### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

**DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** apply where runoff is likely to occur. **DO NOT** apply where weather conditions favor drift from areas treated. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice. Use of this product is prohibited in the following areas where the species are known to exist:

- **Solano Grass:** Solano County, California: the vernal lakes are bounded by the Union Pacific Railroad and Hastings Road to the north, Highway 113 to the east, Highway 12 to the south, and Travis Air Force Base to the west.
- **Wild Rice:** Hays County, Texas

#### NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

#### PHYSICAL OR CHEMICAL HAZARDS

**Combustible.** **DO NOT** use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves including Barrier Laminote or Viton  $\geq$  14 mils
- Shoes plus socks
- Protective eyewear

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas or vicinity where there may be drift. **DO NOT** enter treated areas without protective clothing until sprays have dried.

## RESISTANCE MANAGEMENT

**Tigris PS Clethodim** contains clethodim and is classified in the cyclohexanedione chemical group as a Group I herbicide, ACCase inhibitors. Herbicide resistance is defined as the inherited ability of a plant to survive and reproduce following exposure to a dose of herbicide normally lethal to the wild type. In a plant, resistance may be naturally occurring or induced by such techniques as genetic engineering or selection of variants produced by tissue culture or mutagenesis. Any weed population may contain or develop plants that are naturally resistant to **Tigris PS Clethodim** and other Group I herbicides. Weed species with acquired resistance to Group I herbicides may eventually dominate the weed population if Group I herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **Tigris PS Clethodim** or other Group I herbicides.

To delay herbicide resistance take one or more of the following steps:

- Plant into weed-free fields and keep fields as weed-free, as possible.
- To the extent possible, use a diversified approach toward weed management. Whenever possible, incorporate multiple weed-control practices such as mechanical cultivation, biological management practices, and crop rotation.
- Fields with difficult to control weeds should be rotated to crops that allow the use of herbicides with alternative mechanisms of action or different management practices.
- To the extent possible **DO NOT** allow weed escapes to produce seeds, roots, or tubers. Manage weed seeds at harvest and post-harvest to prevent a buildup of the weed seed-bank.
- Prevent field-to-field and within-field movement of weed seed or vegetative propagules. Thoroughly clean plant residues from equipment before leaving fields.
- Prevent an influx of weeds into the field by managing field borders.
- Identify weeds present in the field through scouting and field history and understand their biology. The weed-control program should consider all of the weeds present.
- Difficult to control weeds may require sequential applications of herbicides with differing mechanisms of action.
- Apply this herbicide at the correct timing and rate needed to control the most difficult weed in the field.
- Use a broad-spectrum soil-applied herbicide with a mechanism of action that differs from this product as a foundation in a weed-control program. **DO NOT** use more than 2 applications of this or any other herbicide with the same mechanism of action within a single growing season unless mixed with an herbicide with another mechanism of action with an overlapping spectrum for the difficult-to-control weeds.
- Report any incidence of non-performance of this product against a particular weed species to your Tigris, LLC retailer, representative or call (252) 235-0043.
- Monitor treated weed populations for loss of field efficacy.
- Scout field(s) before and after application.

Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species.

Contact your local sales representative, extension agent, or certified crop advisor to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. tank mix products so that there are multiple effective mechanisms of action for each target weed.

## TANK MIXES

**NOTICE:** It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Those concerns may include, but are not limited to:

1. Geographic restrictions - all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
2. Crop rotation restrictions;
3. Applicator certification requirements;
4. Worker safety rules (including protective clothing, reentry time, posting);
5. Soil type or soil characteristics (including pH,  $\text{O}_2$ );
6. Maximum dosage or number of applications per year;
7. Rain free period required; or
8. Application timing (including pre-harvest interval).
9. **DO NOT** exceed the total yearly rates.

## PRODUCT INFORMATION

### RESTRICTIONS

- **Long Island, New York:** **DO NOT** apply more than 16 fl. oz. of **Tigris PS Clethodim** (0.25 lb. a.i.) per acre per season.
- **DO NOT** apply if rain is expected within 1 hour of application as control may be unsatisfactory.
- **DO NOT** apply a post-emergence broadleaf herbicide within 1 day following application of **Tigris PS Clethodim** or reduced grass control may result.
- **DO NOT** apply under conditions of stress. Applying **Tigris PS Clethodim** under conditions that do not promote active grass growth will reduce herbicide effectiveness. These conditions include drought, excessive water, extremes in temperature, low humidity and grasses either partially controlled or stunted from prior pesticide applications. Grasses under these kinds of stressful conditions will not absorb and translocate **Tigris PS Clethodim** effectively, and will be less susceptible to herbicide activity.
- **DO NOT** allow **Tigris PS Clethodim** to come in contact with desirable grass crops including corn, rice, sorghum, small grains, or turf, as these and other grass crops will be injured or killed. Minor leaf spotting may occur on treated plants under certain environmental conditions. New foliage is not affected.
- Aerial applications for all tree fruits and tree nuts uses are prohibited.
- **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

### ROTATIONAL RESTRICTION

- **DO NOT** plant rotational crops until 30 days after application of **Tigris PS Clethodim** unless the crop is listed on **Tigris PS Clethodim** label.

## PRECAUTIONS

- Grass crops including corn, rice, sorghum, small grains, or turf, etc. are highly sensitive to **Tigris PS Clethodim**.
- **Tigris PS Clethodim** is a selective post-emergence herbicide for control of annual and perennial grasses.
- **Tigris PS Clethodim** does not control sedges or broadleaf weeds.
- Optimal perennial grass control can be obtained if rhizomes or stolons are cut up by pre-plant tillage practices, (discing, plowing, etc.) to stimulate maximum emergence of grass shoots. Cultural practices, including continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up, result in a very staggered, non-uniform weed emergence. Due to this non-uniform weed emergence, no fewer than 2 **Tigris PS Clethodim** applications per year are specified at the appropriate weed growth stage rate under continuous no-till conditions.
- While all the vegetable crops on this label have been tested for crop safety with **Tigris PS Clethodim**, not all specialty varieties of these crops have been tested. It is advised that, before applying **Tigris PS Clethodim** to specialty varieties of vegetable crops on this label, first apply a small section of the field to evaluate potential injury. Injury symptoms may appear as leaf speckling or stunting.
- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- Tank mixes of **Tigris PS Clethodim** and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs, an additional application of **Tigris PS Clethodim** may be necessary. **AVOID SPRAY DRIFT**.
- Repeated use of **Tigris PS Clethodim** (or similar post-emergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes that are resistant to these products in some grass species.
- If poor performance occurs and cannot be attributed to adverse weather or application conditions, a resistant biotype may be present. This is most likely to occur in fields where other control strategies including crop rotation, mechanical removal, and other classes of herbicides are not used from year to year.

## APPLICATION INFORMATION

### Control Symptoms

Treated grass weeds show a reduction in vigor and growth. Early chlorosis/necrosis of younger plant tissue is followed by a progressive collapse of the remaining foliage. Symptoms will generally be observed in 7 to 14 days after application, depending on grass species treated and environmental conditions.

### Timing of Applications

Apply **Tigris PS Clethodim** post-emergence to actively growing grasses according to rate table instructions. Applications made to grass plants stressed by insufficient moisture, or hot or cold temperatures, or to grass plants exceeding listed growth stages may result in unsatisfactory control. **DO NOT** apply under these conditions.

In arid regions where irrigation is used to supplement limited rainfall, **Tigris PS Clethodim** must be applied as soon as possible, after irrigation (within 7 days). In arid regions, a second application of **Tigris PS Clethodim** will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

Cultivation of treated grasses 7 days prior to or within 7 days after application of **Tigris PS Clethodim** may reduce weed control. **DO NOT APPLY Tigris PS Clethodim** if rainfall is expected within 1 hour, since control may be reduced.

### ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE

CROP	ADJUVANT USE INSTRUCTIONS
Alfalfa, Cotton, Legume Vegetables (succulent & dried) – Crop Group 6, Peanuts (including perennial), Potato, Soybean, Sugar Beet, and Sunflower – Subgroup 20B.	Always use a crop oil concentrate* at 1.0 qt./A by ground or 1% v/v (but not less than 1 pt./A) in the finished spray volume by air. To 2 qts./A of liquid fertilizer (10-34-0, 28%N or 32%N), or an equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added to <b>Tigris PS Clethodim</b> applications, in addition to the specified rate of crop oil concentrate. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.
Artichoke (Globe), Berry (low-growing) – Subgroup 13-076 (except cranberry), Brassica Head & Stem Vegetables – Crop Group 5-16, Brassica Leafy Greens – Subgroup 4-16B, Bushberry – Subgroup 13-078, Caneberry – Subgroup 13-07A, Clover, Cranberry, Flax, Fruit, Fruiting Vegetables – Group 8-10 (except Okra and Tomato), Garden Beet, Herbs – Subgroup 19A, Hops, Leafy Greens – Subgroup 4-16 Leaf Petiole Vegetable – Subgroup 22B, Melon – Subgroup 9A, Mustard Seed, Okra, Onion (Bulb) – Subgroup 3-07A, Onion (Green) – Subgroup 3-07B, Peppermint and Spearmint tops, Pome Fruit – Crop Group 11-10, Radish, Rapeseed – Subgroup 20A (except flax seed), Root Vegetables – Subgroup 1B (except sugar beet), Safflower, Squash/cucumber – Subgroup 9B, Stalk and Stem Vegetable – Subgroup 22A, Stevia (dried leaves), Stone Fruit – Crop Group 12-12, Tuberous and Corm Vegetable – Subgroup 1C, and Tree Nuts – Crop Group 14-12	Always use a crop oil concentrate at 1% v/v in the finished spray volume unless tank mix instructions indicate otherwise. Addition of liquid fertilizer is not advised for these crops.
Non-Bearing Food Crops, Ornamental Plants	Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v). Use of crop oil concentrate is not advised since it may injure flowers and foliage.
Conifer Trees, Fallow Land (and other non-producing agricultural areas), and Non-Crop or Non-Planted Areas	Always use a crop oil concentrate containing at least 15% emulsifier at 1% v/v (but not less than 1 pt./A) in the finished spray volume.

\*Acceptable crop oil concentrates would be those that contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality, and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

## Ground Application

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 5 gals. and a maximum of 40 gals. of spray solution per acre. Under the following conditions a minimum of 10 gals. per acre is required: ultra-narrow row cotton, narrow row soybeans, broadleaf herbicide tank mixes, perennial grasses, volunteer corn, drought or stress conditions, heavy grass pressure or when grasses are at or near maximum height. Failure to use a minimum of 10 gals. per acre under these conditions can result in poor coverage and reduced grass control requiring repeat applications. Spray pressures must reflect a minimum of 30 PSI and a maximum of 60 PSI at the nozzle. **DO NOT** use flood nozzles.

Applications to onions (dry bulbs and green), garlic, and shallots (dry bulbs and green) must be made in a minimum of 20 gals. of spray solution per acre.

## Air Application

Use a minimum of 3 gals. of spray solution per acre unless otherwise directed in this label. Increase spray volumes up to 10 gals. as grass or crop foliage becomes dense. **Restriction:** For onions (dry bulbs and green), garlic, or shallots (dry bulbs and green): When applying by air **DO NOT** exceed 8 fl. oz./A (0.025 lb. a.i./A) in a single application. In California, air applications to onions, garlic or shallots must be made in a minimum of 20 gals. of spray solution per acre. In states other than California, air application to onions, garlic, or shallots must be made in a minimum of 10 gals. of spray solution per acre.

**NOTE: Crop injury may occur when Tigris PS Clethodim is applied to onions, garlic, or shallots with aerial equipment.**

## Spot Treatment

When using hand sprayers or high-volume sprayers utilizing handgunns, mix 1/4 to 1/2% (0.33 oz. to 0.65 oz. per gal) **Tigris PS Clethodim** and treat to wet vegetation, while not allowing runoff of spray solution. For uses requiring crop oil concentrate, include crop oil concentrate at 1% (1.3 oz. per gal) by volume. For uses requiring non-ionic surfactant, include non-ionic surfactant at 1/4% (0.33 oz. per gal) by volume.

**NOTE: If Tigris PS Clethodim is applied as a spot treatment, care must be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.**

## CHEMIGATION – ONION (Dry Bulb and Green) AND GARLIC SPRINKLER IRRIGATION APPLICATION

Only apply **Tigris PS Clethodim** by chemigation to Onions (dry bulb and green) and Garlic Crops.

May be applied to onions and garlic by sprinkler irrigation systems.

Apply **Tigris PS Clethodim** at the high rate specified for annual grasses (16 fl. oz. per acre (0.25 lb. a.i.)) when the grass height is at the low end of the range (application to larger grasses may not provide adequate control). Add a crop oil concentrate containing at least 15% emulsifier at 1 quart per acre.

Apply **Tigris PS Clethodim** in 0.1 to 0.2 acre-inch of water either at the end of a regular irrigation set or as a separate application not associated with a regular irrigation that provides proper distribution and coverage. Application of more than label-specified quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness. Use a metering device to inject the **Tigris PS Clethodim** into the irrigation water at a constant flow. Constant agitation must be maintained in the chemical supply tank during the entire period of herbicide application. Inject the product with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period.

## PRECAUTIONS

- Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set, or hand move.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.
- If you have any questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

## RESTRICTIONS

- **DO NOT apply Tigris PS Clethodim by chemigation in the states of Idaho, Montana, Oregon, and Washington.**
- **DO NOT** apply this product through any irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- **DO NOT** apply by chemigation to any other crop, or to this crop using any other type of irrigation system.
- **DO NOT** apply this product through any other type of irrigation system (i.e., Public Irrigation System).
- **DO NOT** connect to an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.
- A person knowledgeable of the chemigation system and responsible for its operation or under supervision of the responsible person, must shut the system down and make necessary adjustments should the need arise.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, including a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

## MANDATORY SPRAY DRIFT

### Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applications must use  $1/2$  swath displacement upward at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed towards the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

### Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy. For all other ground applications, the nozzle must be no more than 3 feet from the target vegetation.
- Applicators are required to use a medium or coarser droplet size (ASABE S572).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

## SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using a nozzle designed to reduce drift.

### Control Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### BOOM HEIGHT - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

### SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## PRODUCT LIMITATIONS

Optimal perennial grass control can be obtained if rhizomes or stolons are cut up by pre-plant tillage practices, (discing, plowing, etc.) to stimulate maximum emergence of grass shoots. Cultural practices, including continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up, result in a very staggered, non-uniform weed emergence. Due to this non-uniform weed emergence, no fewer than 2 **Tigris PS Clethodim** applications per year are advised at the appropriate weed-growth stage rate under continuous no-till conditions.

Grass crops including corn, rice, sorghum, small grain, or turf, etc. are highly sensitive to **Tigris PS Clethodim**.

While all the vegetable crops on this label have been tested and are resistant to **Tigris PS Clethodim**, not all specialty varieties of these crops have been tested. It is advised that, before applying **Tigris PS Clethodim** to specialty varieties of vegetable crops on this label, crop resistance be investigated first using a small section of the field. It is possible that injury symptoms can occur. Symptoms may appear as leaf speckling or stunting.

Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.

Tank mixes of **Tigris PS Clethodim** and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs, an additional application of **Tigris PS Clethodim** may be necessary. AVOID SPRAY DRIFT.

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PH)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Alfalfa</b> including: Sainfoin Holy Clover Birdsfoot Trefoil <sup>(3)</sup>	15 days before grazing, feeding or harvesting (cutting) for forage or hay	6-16 fl. oz. <sup>(4)</sup> (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(5)</sup>	<p>Refer to tank mix partners for feeding, grazing, and harvesting restrictions.<sup>(6)</sup></p> <p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.25 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz./A (0.25 lb. a.i./A) per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply <b>Tigris PS Clethodim</b> plus 2,4-DB as a tank mix to alfalfa unless the 60 day feeding, grazing, and harvesting restriction on the 2,4-DB label can be observed.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Artichoke (Globe)</b>	5 days	6-8 fl. oz. (0.094-0.25 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Bean, Dry except Soybean [Dried Shelled Pea and Bean (except soybean) Subgroup 6C]</b> including: Bean ( <i>Lupinus</i> spp.) Grain, Sweet, White, White Sweet Bean ( <i>Phaseolus</i> spp.) Field, Kidney, Lima (dry), Navy, Pinto, Tepary Bean ( <i>Vigna</i> spp.) Adzuki Bean, Black-eyed Pea, Catjang, Cowpea, Crowder Pea, Math Bean, Mung Bean, Rice Bean, Southern Pea, Urd Bean, Broad (dry), Chickpea (garbanzo), Guar, Lablab Bean, Lentil	30 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(5)</sup>	<p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.25 lb. a.i./A), or 2 applications at 16 fl. oz./A (0.25 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For Reduced Rate Control for Small Annual Grasses: <b>DO NOT</b> make more than 8 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul> <p>Refer to the <b>DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES</b> table.</p>
<b>Bean, Succulent Shelled [Succulent Shelled Pea and Bean Subgroup 6B]</b> including: Bean ( <i>Phaseolus</i> spp.) Broad Bean (succulent), Lima Bean (green) Bean ( <i>Vigna</i> spp.) Black-eyed Pea, Cowpea, Southern Pea	21 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(5)</sup>	<p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply more than 1 application per acre per year.</li> <li>• For Reduced Rate Control for Small Annual Grasses: <b>DO NOT</b> make more than 2 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year. For repeat applications, make on a minimum of a 14-day interval.</li> </ul> <p>Refer to the <b>DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES</b> table.</p>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

<sup>3</sup> **Tigris PS Clethodim** may be applied to seedling or established alfalfa grown for seed, hay, silage, green crop, or direct grazing.

<sup>4</sup> For weed control in established alfalfa and mint (peppermint and spearmint tops), the minimum use rate is 10 fl. oz./A (0.156 lb. a.i./A).

<sup>5</sup> 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N), or an equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added to **Tigris PS Clethodim** applications, in addition to the specified rate of crop oil concentrate.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PH)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
Beet, Garden	30 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.500 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Berry Low Growing (except Cranberry and Strawberry) [Subgroup 13-07G]</b> including: Bearberry Bilberry Blueberry, Lowbush Cloudberry Lingonberry Muntres Partridgeberry	45 days	4-8 fl. oz. (0.0625-0.125 lb. a.i./A)	Non-ionic surfactant (NIS) at 0.25% v/v.	Verify crop safety to <b>Tigris PS Clethodim</b> on a small area of the crop, at the desired <b>Tigris PS Clethodim</b> rate and with the same non-ionic surfactant (NIS) that will be used on the field. If no crop response is evident 7 days after treatment, <b>Tigris PS Clethodim</b> may be used on the entire field at the rate tested and with the same NIS used in the crop safety test. <b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.500 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Brassica Head &amp; Stem Vegetable [Crop Group 5-16]</b> including: Broccoli Brussels Sprouts Cabbage Cabbage, Chinese (Napa) Cauliflower Cultivars, varieties, and/or hybrids of these commodities.	30 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> apply more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.500 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Brassica Leafy Greens (except Radish Leaves, Turnip Greens and Watercress) [Subgroup 4-16B]</b> including: Arugula Broccoli, Chinese Broccoli Raab Cabbage, Abyssinian Cabbage, Chinese (Bok Choy) Cabbage, Seakale Collards Cress, Garden Cress, Upland Hanover Salad Kale Maca (Leaves) Mizuna Mustard Greens Rape Greens Rocket, Wild Shepherd's Purse Cultivars, varieties, and/or hybrids of these commodities.	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> apply more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Bushberry [Subgroup 13-07B]</b> including: Aronia Berry Blueberry, Highbush Chilean Guava Cranberry, Highbush Currant, Black Currant, Buffalo Currant, Native Currant, Red Elderberry European Barberry Gooseberry Honeysuckle, Edible Huckleberry Jostaberry Juneberry Saskatoon Berry Native Currant Salal Sea Buckthorn Cultivars, varieties, and/or hybrids of these.	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	Apply at the base of the plant where grassy weeds are growing near the ground. <b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply to low growing berries.</li> <li>• <b>DO NOT</b> apply to Bushberry grown for root stock.</li> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply more than 4 applications per acre per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PH)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Caneberry [Subgroup 13-07A]</b> including: Blackberry Loganberry Raspberry, Black Raspberry, Red Raspberry, Wild Cultivars, varieties, and/or hybrids of these.	7 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p>Apply at the base of the plant where grassy weeds are growing near the ground.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> apply more than 4 applications when using 8 fl. oz./A (0.125 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply to low growing berries.</li> <li>• <b>DO NOT</b> apply to Caneberry grown for root stock.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Canola (except flax seed, mustard seed and sesame seed) [Rapeseed Subgroup 20A]*</b> including: Borage Crambe Cuphea Echium Gold of Pleasure (Camelina) Hare's Ear Mustard Lesquerella Lunaria Meadowfoam Milkweed Oil Radish Poppy Seed Rapeseed (Canola) Sweet Rocket Cultivars, varieties, and/or hybrids of these. *Not for use in California.	70 days	4-6 fl. oz. (0.0625-0.094 lb. a.i./A)	1% v/v in the finished spray volume.	<p>Verify crop safety to <b>Tigris PS Clethodim</b> on a small area of the crop, at the desired <b>Tigris PS Clethodim</b> rate and with the same non-ionic surfactant (NIS) that will be used on the field. If no crop response is evident 7 days after treatment, <b>Tigris PS Clethodim</b> may be used on the entire field at the rate tested and with the same NIS used in the crop safety test.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply after crop has begun bolting. Crop injury may occur when <b>Tigris PS Clethodim</b> is applied during the bloom period.</li> <li>• <b>DO NOT</b> apply more than 6 fl. oz./A (0.094 lb. a.i./A) in a single application.</li> <li>• For Annual and Perennial Grasses: <b>DO NOT</b> make more than 1 application at 6 fl. oz./A (0.094 lb. a.i./A) in a year.</li> <li>• For Reduced Rate Control for Small Annual Grasses: <b>DO NOT</b> make more than 1 application at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> </ul> <p>Refer to the <b>DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES</b> table.</p>
<b>Carrot</b>	30 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Clover</b>	15 days before grazing, feeding, or harvesting (cutting) for forage or hay	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1% v/v in the finished spray volume.	<p>For use on clover grown in the states of Idaho, Oregon, and Washington only.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 2 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 1 application at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per acre per year.</li> <li>• <b>DO NOT</b> exceed 16 fl. oz./A (0.25 lb. a.i./A) in a year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Corn, Field</b> For burndown of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting field corn. See Directions for Use in Roundup Ready Field Corn (Burndown) table.	90 days	3 fl. oz. (0.047 lb. a.i./A)	<b>DO NOT</b> use COC or MSO with <b>Tigris PS Clethodim</b> in this use pattern.	To control the existing stand, replant no sooner than 6 days after application. <b>Restrictions:</b> <ul style="list-style-type: none"><li>• <b>DO NOT</b> apply more than 3 fl. oz./A (0.047 lb. a.i./A) per application.</li><li>• <b>DO NOT</b> make more than 1 application per year.</li><li>• <b>DO NOT</b> apply more than 3 fl. oz./A (0.047 lb. a.i./A) per year.</li></ul>
<b>Cotton (including cotton grown for seed) [Cottonseed Subgroup 20C]</b> including: Cottonseed Cultivars, varieties, and/or hybrids of these.	60 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(3)</sup>	The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. <b>Restrictions:</b> <ul style="list-style-type: none"><li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li><li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 6 fl. oz. (0.25 lb. a.i./A) per acre per year.</li><li>• <b>DO NOT</b> apply more than 32 fl. oz. (0.5 lb. a.i./A) per year.</li><li>• <b>DO NOT</b> graze treated fields or feed treated forage or hay to livestock.</li><li>• For repeat applications, make on a minimum of a 14-day interval.</li></ul>
<b>Cranberry</b>	30 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"><li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li><li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li><li>• <b>DO NOT</b> apply more than 32 fl. oz. (0.5 lb. a.i./A) per year.</li><li>• <b>DO NOT</b> apply between the "hook" stage and full fruit set.</li><li>• For repeat applications, make on a minimum of a 14-day interval.</li></ul>
<b>Fallow Land</b> <b>Conifer Trees</b> (and other non-producing agricultural areas) <b>Non-Crop or Non-Planted Areas</b>	N/A	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1% v/v (but not less than 1 pt./A) in the finished spray volume using a crop oil concentrate containing at least 15% emulsifier.	<b>Restrictions:</b> <ul style="list-style-type: none"><li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li><li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 6 fl. oz. (0.25 lb. a.i./A) per acre per year.</li><li>• <b>DO NOT</b> apply more than 32 fl. oz. (0.5 lb. a.i./A) per year.</li><li>• <b>DO NOT</b> plant any crop for 30 days after application unless clethodim is registered for use in that crop.</li><li>• For repeat applications, make on a minimum of a 14-day interval.</li></ul>
<b>Flax</b> <sup>*</sup> <sup>*Not for use in California.</sup>	60 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	Apply prior to bloom. Crop injury may occur when <b>Tigris PS Clethodim</b> is applied during the bloom period. <b>Restrictions:</b> <ul style="list-style-type: none"><li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li><li>• <b>DO NOT</b> make more than 2 applications per acre per year.</li><li>• <b>DO NOT</b> exceed 16 fl. oz./A (0.25 lb. a.i./A) in a year.</li><li>• For Reduced Rate Control for Small Annual Grasses: <b>DO NOT</b> make more than 4 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li><li>• For repeat applications, make on a minimum of a 14-day interval.</li></ul> Refer to the <b>DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES</b> table.

N/A = Not Applicable

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

<sup>3</sup> 1 to 2 qt./A of liquid fertilizer (10-34-0, 28%N, or 52%N), or an equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added to **Tigris PS Clethodim** applications, in addition to the specified rate of crop oil concentrate.

(continued)

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PH)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Fruiting Vegetable (except Okra and Tomato) [Crop Group 8-10]</b> including: African Eggplant Bush Tomato Bell Pepper Cocona Currant Tomato Eggplant Garden Huckleberry Goji Berry Groundcherry Martynia Maronjilla Pea Eggplant Pepino Non-bell Pepper Roselle Scarlet Eggplant Sunberry Tomatillo Tree Tomato Cultivars, varieties, and/or hybrids of these.	20 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use Tigris PS Clethodim on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Herbs [Subgroup 19A]</b> including: Angelica Balm Basil Borage Burnet Camomile Catnip Chervil (Dried) Chive Chive, Chinese Clary Coriander (Leaf) Costmary Cilantro (Leaf) Curry (Leaf) Dill (Dillweed) Horehound Hyssop Lavender Lemongrass Lovage (Leaf) Marigold Marjoram ( <i>Origanum</i> spp.) Nasturtium Parsley (Dried) Pennyroyal Rosemary Rue Sage Savory, Summer and Winter Sweet Bay Tansy Tarragon Thyme Wintergreen Woodruff Wormwood	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Tigris PS Clethodim</b> has not been tested on all herbs, and herb varieties. It is the responsibility of the user to test <b>Tigris PS Clethodim</b> on a small portion of the crop to be treated before treating the entire field.</p> <p>Crop resistance must be verified to <b>Tigris PS Clethodim</b> on a small area of the herb crop, at the desired <b>Tigris PS Clethodim</b> rate and with the same crop oil concentrate that will be used on the herb field. If no crop response is evident 7 days after treatment, <b>Tigris PS Clethodim</b> may be used on the entire field at the rate tested and with the same crop oil used in the resistance test.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per acre per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Hops</b>	21 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per acre per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Leaf Petioles Vegetables [Subgroup 22B]*</b> including: Cardoon Celery Celuce Chinese Celery Fuki Fennel, Florence (Finocchio) Rhubarb Udo Zukki Swiss Chard Cultivars, varieties, and/or hybrids of these commodities. <small>*Not for use in California.</small>	30 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>DO NOT make more than 4 applications per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per acre.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Leafy Greens [Subgroup 4-16A]*</b> including: Amaranth Chinese Spinach Leafy Amaranth Tampala Arugula (Roquette) Chervil Chrysanthemum, Edible-leaved Chrysanthemum, Garland Corn Salad Cress Garden Upland (Yellow Rock and Winter) Dandelion Dock (Sorrel) Endive (Escarole) Lettuce, Head and Leaf Orach Parsley Purslane Garden Winter Radicchio (Red Chicory) Spinach New Zealand Vine (Indian and Malabar) <small>*Not for use in California.</small>	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>DO NOT make more than 4 applications per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per acre.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> DO NOT use Tigris PS Clethodim on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE section for further information.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Legume Vegetables, Edible Podded [Subgroup 6A]</b> including: Bean ( <i>Phaseolus</i> spp.) Runner, Snap, Wax Bean ( <i>Vigna</i> spp.) Asparagus, Chinese Longbean, Moth, Yardlong, Jackbean Pea ( <i>Pisum</i> spp.) Dwarf, Edible-pod, Snow, Sugar Snap Pigeon Sword Bean	21 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt.) by air. <sup>(5)</sup>	For peas, apply before bloom, but no later than 21 days before harvest. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. <b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>DO NOT apply more than 1 application per acre per year.</li> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) per year.</li> <li>For Reduced Rate Control for Small Annual Grasses: DO NOT make more than 2 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul> Refer to the <b>DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES</b> table.
<b>Melon [Subgroup 9A]</b> including: Citron Melon Melon Muskmelon (including Cantaloupe, Honeydew) Watermelon	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>DO NOT make more than 4 applications per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Mint (Peppermint &amp; Spearmint Tops)</b>	21 days	6-16 fl. oz. <sup>(4)</sup> (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt.) by air.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 16 fl. oz./A (0.25 lb. a.i./A) in a single application.</li> <li>DO NOT make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Mustard Seed *</b> <small>*Not for use in California.</small>	75 days	4-6 fl. oz. (0.0625-0.094 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply after crop has begun bolting. Crop injury may occur when <b>Tigris PS Clethodim</b> is applied during the bloom period.</li> <li>DO NOT apply more than 6 fl. oz./A (0.094 lb. a.i./A) in a single application.</li> <li>DO NOT make more than 2 applications per acre per year.</li> <li>DO NOT exceed 12 fl. oz./A (0.188 lb. a.i./A) in a year.</li> <li>For Reduced Rate Control for Small Annual Grasses: DO NOT make more than 3 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul> Refer to the <b>DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES</b> table.
<b>Okra</b>	3 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>DO NOT make more than 4 applications per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> DO NOT use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

<sup>4</sup> For weed control in established alfalfa and mint (peppermint and spearmint tops), the minimum use rate is 10 fl. oz./A (0.156 lb. a.i./A).

<sup>5</sup> 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N), or an equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added to **Tigris PS Clethodim** applications, in addition to the specified rate of crop oil concentrate.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Bulb Onion [Subgroup 3-07A]</b> including: <b>Onion (Dry Bulb Only)</b> Including: Daylily, Bulb Fritillaria, Bulb <b>Garlic, Bulb</b> Garlic, Great-headed, Bulb Garlic, Serpent, Bulb Lily, Bulb Onion, Bulb Onion, Chinese, Bulb Onion, Pearl Onion, Potato, Bulb <b>Shallot (Dry Bulb Only)</b> Cultivars, varieties, and/or hybrids of these.	45 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1% v/v in the finished spray volume.	Minimum of 20 gals./A spray volume by ground in entire U.S. Minimum of 20 gals./A spray volume by air in California. In states other than California, air application to onions, garlic or shallots must be made in a minimum of 10 gals./A. <b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li> <li>DO NOT make more than 2 applications per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>When applying by air, DO NOT exceed 8 fl. oz./A (0.125 lb. a.i./A) per application and 16 fl. oz./A (0.25 lb. a.i./A) per year.</li> <li>If <b>Tigris PS Clethodim</b> is applied as a spot treatment to onion or garlic, DO NOT exceed the maximum rate allowed on a "per acre" basis.</li> <li>In California, DO NOT apply <b>Tigris PS Clethodim</b> to garlic or onion until the crop has at least 2 full leaves. Use a 14-day spray interval between the application of <b>Tigris PS Clethodim</b> and liquid nitrogen or other herbicide applications. Injury to crop may occur when shorter intervals are observed.</li> <li>In California, for air applications to onion, DO NOT exceed 2 applications per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul> <b>For Garlic or Shallot:</b> <ul style="list-style-type: none"> <li>When applying by ground, DO NOT exceed 8 fl. oz./A (0.125 lb. a.i./A) per application or 16 fl. oz./A (0.25 lb. a.i./A) per year.</li> <li>DO NOT exceed 2 applications per year.</li> </ul> <b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>DO NOT make more than 4 applications per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>For air applications to onion, DO NOT exceed 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>In California for air applications to onion, DO NOT exceed 2 applications per year.</li> <li>If <b>Tigris PS Clethodim</b> is applied as a spot treatment to onion or garlic, DO NOT exceed the maximum rate allowed on a "per acre" basis.</li> <li>In California, DO NOT apply <b>Tigris PS Clethodim</b> to onion or garlic until the crop has at least 2 full leaves. Use a 14-day spray interval between the application of <b>Tigris PS Clethodim</b> and liquid nitrogen or other herbicide applications. Injury to crop may occur when shorter intervals are observed.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Onion, Green [Subgroup 3-07B]*</b> including: Chive (Fresh Leaves) Chive, Chinese (Fresh Leaves) Elegans Hosta Fritillaria (Leaves) Kurpat Lady's Leek Leek Leek, Wild Onion, Beltsville (Bunching) Onion Fresh) Onion, Green Onion, Macrostemon Onion, Tree (Tops) Onion, Welsh (Tops) <b>Shallot (Fresh Leaves)</b> Cultivars, varieties, and/or hybrids of these commodities. *Not for use in California.	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	

<sup>1</sup> DO NOT use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

(continued)

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Ornamentals</b>	N/A	6-16 fl. oz. (0.094-0.25 lb. a.i.)	Use of Crop Oil Concentrate is not advised since it may injure flowers and foliage.  See Special Use Instructions.	Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).
<b>Non-Bearing Food Crops</b>	N/A	6-8 fl. oz. (0.094-0.125 lb. a.i.)		<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>For Ornamentals, <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) in a single application.</li> <li>For Non-Bearing Food Crops, <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li><b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li><b>DO NOT</b> apply more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per year.</li> <li>Sugar maples cannot be tapped for syrup within 1 year of <b>Tigris PS Clethodim</b> application.</li> <li>If applying as a spot treatment to non-bearing food crops, <b>DO NOT</b> exceed the maximum rate allowed on a "per acre" basis.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Pea, Dry Shelled [Dry Shelled Pea and Bean (except Soybean) Subgroup 6C] including: Pea (<i>Pisum</i> spp.) Field, Pigeon</b>	30 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air <sup>(3)</sup>	<p>Apply before bloom, but not later than 30 days prior to harvest. Applications of <b>Tigris PS Clethodim</b> to peas during the bloom period could result in severe crop injury, including loss of yield and delayed maturity.</p> <p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li><b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li><b>DO NOT</b> apply more than 1 application per acre per year.</li> <li><b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per year.</li> <li>For Reduced Rate Control for Small Annual Grasses: <b>DO NOT</b> make more than 2 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Pea, Succulent [Succulent Shelled Pea and Bean Subgroup 6B] including: Pea (<i>Pisum</i> spp.) English Pea, Garden Pea, Green Pea Pigeon Pea</b>	21 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air <sup>(3)</sup>	<p>Apply before bloom, but not later than 21 days prior to harvest. Applications of <b>Tigris PS Clethodim</b> to peas during the bloom period could result in severe crop injury, including loss of yield and delayed maturity.</p> <p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li><b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li><b>DO NOT</b> apply more than 1 application per acre per year.</li> <li><b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per year.</li> <li>For Reduced Rate Control for Small Annual Grasses: <b>DO NOT</b> make more than 2 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

N/A = Not Applicable

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

<sup>3</sup> 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N), or an equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added to **Tigris PS Clethodim** applications, in addition to the specified rate of crop oil concentrate.

(continued)

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Peanut (including Perennial)</b>	40 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(3)</sup>	<p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Pome Fruit [Crop Group II-10]</b> including: Apple Azarole Crabapple Loquat Mayhaw Hook & Arn., and <i>C. rufa</i> L. Sarg. Medlar Pear Pear, Asian <i>Pseudocydonia sinensis</i> Quince Quince, Chinese Quince, Japanese Tejocote Cultivars, varieties, and/or hybrids of these.	14 days	4-8 fl. oz. (0.0625-0.125 lb. a.i./A)	Non-ionic surfactant (NIS) at 0.25% v/v	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Potato</b>	30 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(3)</sup>	<p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Radish</b> <b>Radish Leaves</b>	15 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 2 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per acre in a year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use Tigris PS Clethodim on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

<sup>3</sup> 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N), or an equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added to Tigris PS Clethodim applications, in addition to the specified rate of crop oil concentrate.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PH)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Root Vegetable (except Sugar Beet &amp; Radish) [Subgroup 1B]</b> including: Burdock, Edible Celeriac Chevrol, Turnip Rooted Chicory Ginseng Horseradish Parsley, Turnip Rooted Parsnip Radish, Oriental Rutabaga Salsify Salsify, Black Salsify, Spanish Skirret Turnip	30 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Safflower</b>	70 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Sesame</b>	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply during flowering.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Soybean</b>	60 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(5)</sup>	<p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A) or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> graze treated fields or feed treated forage or hay to livestock.</li> <li>• For Reduced Rate Control for Small Annual Grasses: <b>DO NOT</b> make more than 8 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul> <p>Refer to the <b>DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES</b> table.</p>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

5 1 to 2 qt./A of liquid fertilizer (10-34-0, 28%N, or 52%N) or an equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added to **Tigris PS Clethodim** applications, in addition to the specified rate of crop oil concentrate.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Squash/Cucumber [Subgroup 9B]</b> including: Chayote (Fruit) Chinese Waxgourd (Chinese Preserving Melon) Cucumber Gherkin Gourd, Edible (Includes Hyatan, Cucuzza, Hechima, Chinese Okra) <i>Momordica</i> spp. (Includes Balsam Apple, Balsam Pear, Bittermelon, Chinese Cucumber) Pumpkin Squash, Summer Squash, Winter (Includes Butternut Squash, Calabaza, Hubbard Squash, Acorn Squash, Spaghetti Squash)	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Stalk and Stem Vegetable [Subgroup 22A]<sup>1</sup></b> Agave Aloe Vera Asparagus Bamboo Shoots Celtuce Fennel, Florence (Fresh Leaves and Stalk) Fern, Fiddlehead (Edible) Kale, Sea Kohlrabi Palm Hearts Prickly Pear (Pads) Prickly Pear, Texas (Pads) Cultivars, varieties, and/or hybrids of these commodities.  *Not for use in California.	1 day	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 2 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Stevia, Dried Leaves</b>	14 days	4-8 fl. oz. (0.0625-0.125 lb. a.i./A)	Non-ionic Surfactant (NIS) at 0.25% v/v.	<p><b>Tigris PS Clethodim</b> has not been tested on all varieties. It is the responsibility of the user to test <b>Tigris PS Clethodim</b> on a small portion of the crop to be treated before treating the entire field.</p> <p>Verify crop safety to <b>Tigris PS Clethodim</b> on a small area of the crop, at the desired <b>Tigris PS Clethodim</b> rate and with the same non-ionic surfactant (NIS) that will be used on the field. If no crop response is evident 7 days after treatment, <b>Tigris PS Clethodim</b> may be used on the entire field at the rate tested and with the same NIS used in the crop safety test.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> DO NOT use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

(continued)

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Stone Fruit [Crop Group 12-12]</b> including: Apricot Apricot, Japanese Capulin Cherry, Black Cherry, Nanking Cherry, Sweet Cherry, Tart Jujube, Chinese Nectarine Peach Plum Plum, American Plum, Beach Plum, Canada Plum, Cherry Plum, Chickasaw Plum, Damson Plum, Japanese Plum, Klamath Plum, Prune Plumcot Sloe Cultivars, varieties, and/or hybrids of these.	14 days	4-8 fl. oz. (0.0625-0.125 lb. a.i./A)	Non-ionic Surfactant (NIS) at 0.25% v/v.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.</li> <li>DO NOT make more than 4 applications per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Strawberry</b>	4 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>DO NOT make more than 4 applications per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Sugar Beet</b>	40 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(5)</sup>	The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. <b>Restrictions:</b> <ul style="list-style-type: none"> <li>DO NOT apply more than 16 fl. oz./A (0.25 lb. a.i./A) in a single application.</li> <li>DO NOT make more than 4 applications at the lowest rate, 6-8 fl. oz./A (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz./A (0.25 lb. a.i./A), per acre per year.</li> <li>DO NOT apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>For Reduced Rate Control for Small Annual Grasses: DO NOT make more than 8 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> <li>For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> DO NOT use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

5 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N), or an equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added to **Tigris PS Clethodim** applications, in addition to the specified rate of crop oil concentrate.

Refer to the **DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES** table.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Sunflower [Subgroup 20B]</b> including: Calendula Castor oil Plant Chinese Tallowtree Euphorbia Evening Primrose Jojoba Niger Seed Rose Hip Stokes Aster Tallowwood Tea Oil Plant Vernonia Cultivars, varieties, and/or hybrids of these.	70 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. <sup>(3)</sup>	The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn. Verify crop safety to <b>Tigris PS Clethodim</b> on a small area of the crop, at the desired <b>Tigris PS Clethodim</b> rate and with the same crop oil concentrate that will be used on the field. If no crop response is evident 7 days after treatment, <b>Tigris PS Clethodim</b> may be used on the entire field at the rate tested and with the same crop oil used in the crop safety test.
<b>Tomato</b>	20 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For Reduced Rate Control for Small Annual Grasses: <b>DO NOT</b> make more than 8 applications at 4 fl. oz./A (0.0625 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul> <b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.125 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

<sup>5</sup> 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N), or an equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added to **Tigris PS Clethodim** applications, in addition to the specified rate of crop oil concentrate.

*(continued)*

## CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PH)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Tree Nuts [Crop Group 14-12]</b> including: African Nut-tree Almond Beechnut Brazil Nut Brazilian Pine Bunya Bur Oak Butternut Cajou Nut Condlenut Cashew Chestnut Chinquapin Coconut Coquito Nut Diko Nut Ginkgo Guiana Chestnut Hazelnut (filbert) Heartnut Hickory Nut Japanese Horse-Chestnut Macadamia Nut Mongongo Nut Monkey-pot Monkey Puzzle Nut Okari Nut Pochira Nut Peach Palm Nut Pecan Pequi Pili Nut Pine Nut Pistachio Sapucaia Nut Tropical Almond Walnut, Black Walnut, English Yellowhorn Cultivars, varieties, and/or hybrids of these.	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

*(continued)*

**CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM (cont'd)**

Crops <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre <sup>(2)</sup>	Special Use Instructions
<b>Tuberous and Corm Vegetables [Subgroup IC] (except Potato)</b> including: Arracacha Arrowroot Artichoke (Chinese, Jerusalem) Canna, Edible Cassava (Bitter, Sweet) Choyote (Root) Chufo Dasheen Ginger Leren Sweet Potato Tonier Tumeric Yam Bean Yam, True	30 days	6-16 fl. oz. (0.094-0.25 lb. a.i./A)	1% v/v in the finished spray volume.	<p>The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals, and volunteer corn.</p> <p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications at the lowest rate, 6-8 fl. oz. (0.094-0.25 lb. a.i./A), or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./A), per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Turnip Greens</b>	14 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume.	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>
<b>Watercress<sup>(3)</sup></b> *Not for use in California.	30 days	6-8 fl. oz. (0.094-0.125 lb. a.i./A)	1% v/v in the finished spray volume	<p><b>Restrictions:</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 8 fl. oz./A (0.125 lb. a.i./A) in a single application.</li> <li>• <b>DO NOT</b> make more than 4 applications per acre per year.</li> <li>• <b>DO NOT</b> apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply when watercress is under flood conditions.</li> <li>• <b>DO NOT</b> apply <b>Tigris PS Clethodim</b> when water is in the field and hold water for at least 24 hours after an application.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>

<sup>1</sup> **DO NOT** use **Tigris PS Clethodim** on vegetable crops being grown for seed production unless specific use directions are provided.

<sup>2</sup> Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the **ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE** section for further information.

**IMPORTANT**

Plant resistance to **Tigris PS Clethodim** at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is advised that the user determine if the herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of **Tigris PS Clethodim** have investigated the safety factor to plants not listed on the label.

**DIRECTIONS FOR NON-BEARING FOOD CROPS****TIGRIS PS CLETHODIM MUST NOT BE APPLIED TO NON-BEARING FRUIT OR NUT CROPS WHICH ARE GROWN FOR ROOT STOCK.**

Crop injury to non-bearing fruit and nut crops can occur if **Tigris PS Clethodim** is improperly applied. **Tigris PS Clethodim** must not be applied directly over the top of these plant types. Instead, spray must be directed at the base of the plant where grassy weeds are growing near the ground.

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least 1 year following **Tigris PS Clethodim** application.

COMMON NAME	SCIENTIFIC NAME
Apples	<i>Malus</i> spp.
Berries	<i>Vaccinium</i> spp. <i>Rubus</i> spp.
Cherry, Sweet	<i>Prunus avium</i>
Citrus Fruits	<i>Citrus</i> spp.
Grapes	<i>Vitis</i> spp.
Olives	<i>Olea</i> spp.
Peach	<i>Prunus persica</i>
Pears	<i>Pyrus communis</i>
Prunes	<i>Prunus</i> spp.
Stone Fruits	<i>Prunus</i> spp.
Strawberries	<i>Fragaria</i> spp.
<b>Tree Nuts</b>	
Almond	<i>Prunus triloba</i>
Filbert	<i>Corylus maxima</i>
Pecan	<i>Carya illinoiensis</i>
Pistachio	<i>Pistacia vera</i>
Walnut	<i>Juglans</i> spp.

**Use Rates Per Acre:** 6-8 fl. oz. (0.094-0.125 lb. a.i./A)

**Use Instructions:**

- Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).
- Use of crop oil concentrate is not advised since it may injure flowers and foliage.
- Sugar maples cannot be tapped for syrup within 1 year of **Tigris PS Clethodim** application.

**Restrictions:**

- If **Tigris PS Clethodim** is applied as a spot treatment to non-bearing fruit and nut crops, **DO NOT** exceed the maximum rate allowed on a "per acre" basis.
- **Tigris PS Clethodim** must not be applied to non-bearing fruit or nut crops which are grown for root stock.
- **DO NOT** apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application.
- **DO NOT** make more than 4 applications per acre per year.
- **DO NOT** apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.
- For repeat applications, make on a minimum of a 14-day interval.

### CONIFER TREES

**Tigris PS Clethodim** can be used to control labeled grasses in Christmas tree farms, conifer nurseries, and conifer plantations (but not in forests).

COMMON NAME	SCIENTIFIC NAME
Arborvitae, American	<i>Thuja occidentalis</i>
Cedars	<i>Cedrus</i> spp.
Cypress	<i>Taxodium</i> spp.
Fir, Douglas	<i>Pseudotsuga menziesii</i>
Firs	<i>Abies</i> spp.
Hemlock, Canadian/Eastern	<i>Tsuga Canadensis</i>
Hemlock, Western	<i>Tsuga heterophylla</i>
Pines	<i>Pinus</i> spp.
Spruces	<i>Picea</i> spp.
Yew	<i>Taxus</i> spp.

**Use Rates per Acre:** 6-16 fl. oz. (0.094-0.125 lb. a.i./A)

**Use Instructions:**

- Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).
- **DO NOT** apply more than 16 fl. oz./A (0.25 lb. a.i.) per application.
- **DO NOT** make more than 2 applications per acre per year at the 16 fl. oz. (0.25 lb. a.i./A) rate. **DO NOT** make more than 5 applications per acre per year at the 6 fl. oz. (0.094 lb. a.i./A) rate.
- **DO NOT** apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.
- For repeat applications, make on a minimum of a 14-day interval.

### DIRECTIONS FOR NON-CROP OR NON-PLANTED AREAS

The following areas are considered non-crop or non-planted areas: rights-of-way including railroads, highways, roads, dividers, medians, pipelines, public utility lines, pumping stations, transformer stations and substations. Around airports, electric utilities, commercial buildings, manufacturing plants, storage yards, rail yards, fence lines, parkways, and post-harvest croplands. Also beneath greenhouse benches and around golf courses.

GRASS SUPPRESSION IN NON-CROP AREAS WITH TIGRIS PS CLETHODIM			
GRASS SPECIES	WEED STAGE	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE
Annual and perennial grasses that exceed height claimed for control on height charts below.	Up to and including grasses in the seed head stage.	12 (0.188 lb. a.i.)	16 (0.25 lb. a.i.)
<b>Restrictions:</b>			
<ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply as part of a tank mix when applying <b>Tigris PS Clethodim</b> for grass suppression.</li> <li>• Add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.</li> <li>• <b>DO NOT</b> apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.</li> <li>• <b>DO NOT</b> apply more than 64 fl. oz./A (1 lb. a.i./A) per year.</li> <li>• <b>DO NOT</b> apply more than 4 applications per year.</li> <li>• For repeat applications, make on a minimum of a 14-day interval.</li> </ul>			

**DIRECTIONS FOR ANNUAL GRASSES**  
**(EXCEPT FOR IN ESTABLISHED ALFALFA AND PEPPERMINT AND SPEARMINT TOPS)**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

**Restrictions:**

- DO NOT exceed the maximum rate per application listed in the table **CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM**.
- DO NOT exceed the maximum number of applications per year listed in the table **CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM**.
- DO NOT exceed the maximum yearly rate listed in the table **CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM**.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (Inches)	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE (†)
Barnyardgrass	<i>Echinochloa crus-galli</i>	2 to 8	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Broodleaf Signalgrass	<i>Bracharia platyphylla</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Brome				
California	<i>Bromus carinatus</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Cheat	<i>Bromus secalinus</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Downy	<i>Bromus tectorum</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Rippgrat	<i>Bromus diandrus</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Crabgrass				
Hairy	<i>Digitaria adscendens</i>	2 to 6**	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Large	<i>Digitaria sanguinalis</i>	2 to 6**	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Southern	<i>Digitaria ciliaris</i>	2 to 6**	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Fall Panicum	<i>Panicum dichotomiflorum</i>	2 to 8	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Green	<i>Setaria viridis</i>	2 to 8	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Yellow	<i>Setaria glauca</i>	2 to 8	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)

\* Generally occurs between 3-leaf stage and tillering.

\*\* Length of lateral growth.

(†) Rates higher than 8 fl. oz./A (0.125 fl. a.i./A) may be applied in certain geographic areas, cropping situations, or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 8 to 16 fl. oz./A (0.125 to 0.25 lb. a.i./A) may be applied.

(continued)

**DIRECTIONS FOR ANNUAL GRASSES (cont'd)**

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE (1)
Goosegrass	<i>Eleusine indica</i>	2 to 6**	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Itchgrass	<i>Rottboellia cochinchinensis</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Junglerice	<i>Echinochloa colona</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Red Rice	<i>Oryza sativa</i>	1 to 3	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Ryegrass				
Hardy	<i>Lolium remotum</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Italian	<i>Lolium multiflorum</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Shattercane	<i>Sorghum bicolor</i>	6 to 18	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Sprangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Mexican	<i>Leptochloa uninervia</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Red	<i>Leptochloa filiformis</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Texas Panicum	<i>Panicum texanum</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Volunteer Cereals <sup>(2)</sup>				
Barley	<i>Hordeum vulgare</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Oats	<i>Avena sativa</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Rye	<i>Secale cereal</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Wheat	<i>Triticum aestivum</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)

\* Generally occurs between 3-leaf stage and tillering.

\*\* Length of lateral growth.

(1) Rates higher than 8 fl. oz./A (0.125 fl. a.i./A) may be applied in certain geographic areas, cropping situations, or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 8 to 16 fl. oz./A (0.125 to 0.25 lb. a.i./A) may be applied.

(2) When a cereal grain crop (including wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum **Tigris PS Clethodim** use rate for control is 8 fl. oz./A (0.125 lb. a.i./A).

(continued)

**DIRECTIONS FOR ANNUAL GRASSES (cont'd)**

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (inches)	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE (1)
Volunteer Corn <sup>(3)</sup>	<i>Zea mays</i>	4 to 12	4 (0.0625 lb. a.i.)	6 (0.094 lb. a.i.)
Volunteer Corn <sup>(3)</sup>	<i>Zea mays</i>	12 to 24	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Volunteer Corn (SR) <sup>(4)</sup>	<i>Zea mays</i>	4 to 12	8 (suppression only) (0.125 lb. a.i.)	
Volunteer Grain Sorghum	<i>Sorghum bicolor</i>	8 to 12	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Wild Oats	<i>Avena fatua</i>	2 to 6	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Witchgrass	<i>Panicum capillare</i>	2 to 8	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)

\* Generally occurs between 3-leaf stage and tillering.

(1) Rates higher than 8 fl. oz./A (0.125 fl. oz./A) may be applied in certain geographic areas, cropping situations, or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 8 to 16 fl. oz./A (0.125 to 0.25 lb. a.i./A) may be applied.

(3) Includes Roundup Ready<sup>®</sup>, LibertyLink<sup>®</sup>, and IMI-CORN<sup>®</sup> volunteer corn.

(4) Sethoxydym resistant volunteer corn.

**DIRECTIONS FOR ANNUAL & PERENNIAL GRASS CONTROL IN ESTABLISHED ALFALFA AND PEPPERMINT AND SPEARMINT TOPS WITH TIGRIS PS CLETHODIM**

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE
Annual and Perennial Grasses Listed in Grass Table	See Table	10 (0.156 lb. a.i.)	16 (0.25 lb. a.i.)

**Mowing:** The best control of annual grasses can be achieved by applying **Tigris PS Clethodim** before grass weeds are mowed. Once grass is mowed it becomes tougher to control, as much of the available leaf surface has been removed. In areas without a killing frost, some annuals can over-winter after having been mowed multiple times. These grasses form large crowns and may contain many viable buds. These grasses, even though they may be an annual grass, may require repeated applications of **Tigris PS Clethodim** for partial or complete control.

**Irrigated Alfalfa and Mint (peppermint and spearmint tops):** Irrigation practices can be very critical to the successful use of **Tigris PS Clethodim** in established alfalfa and mint (peppermint and spearmint tops) and may be necessary to initiate active growth of the weeds prior to application. Generally applications 2 to 4 days after irrigation are most effective. Irrigation made shortly after application (2 days) can be effective, but more consistent grass control occurs when the irrigation is made before the application.

**Aerial Application:** Apply **Tigris PS Clethodim** in a minimum of 10 GPa in established alfalfa and mint (peppermint and spearmint tops) when applying by air.

**Annual Grass Control:** Apply **Tigris PS Clethodim** at the grass sizes indicated in the **ANNUAL GRASSES** table and rates indicated. If a grass has been cut, apply **Tigris PS Clethodim** after active growth has resumed and regrowth has reached the minimum height and before it reaches the maximum height indicated. Apply before the alfalfa/mint (peppermint and spearmint tops) canopy covers the grasses and interferes with the spray coverage. Some annual grasses are spring- and summer-germinating plants, while others are fall-germinating plants, and the time they are actively growing and most susceptible to **Tigris PS Clethodim** may vary from region to region. Also some annuals germinate over an extended period of time, and because control of small grasses is desired, applications after each weed flush may be required. As a general rule spray spring- and summer-germinating grasses as early in the season as possible, after initial green-up. Spray fall-germinating weeds in the fall soon after they begin growing but before any damage is done due to frost. Late fall applications may be less effective due to environmental conditions, including frost, slower plant growth, or the onset of flowering.

**Perennial Grass Control:** **Tigris PS Clethodim** effectively controls perennial grasses including Bermudagrass, Johnsongrass, quackgrass, wirestem muhly, tall fescue, foxtail barley and orchardgrass. Due in part to lack of tillage, perennial grasses are more difficult to control in a perennial crop including established alfalfa or mint (peppermint or spearmint tops). A program of repeated applications is usually necessary for best results. The best way to control perennial grasses is to do so in the year of stand establishment before rhizomes and stolons become large and difficult to kill.

Use the high rate under heavy grass pressure and/or when grasses are at or near maximum height.

Always add a crop oil concentrate at 1 qt./A by ground or 1% v/v (but not less than 1 pt./A) to the finished spray volume by air.

**DIRECTIONS FOR ANNUAL BLUEGRASS CONTROL WITH TIGRIS PS CLETHODIM**

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE
Annual Bluegrass ( <i>Poa annua</i> )	to 4-leaf	6* (0.094 lb. a.i.)	16 (0.25 lb. a.i.)

• Apply under favorable soil moisture and humidity, which exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).  
 • Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.  
 • Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.  
 • Always add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.  
 \*Use a minimum of 10 fl. oz./A (0.156 lb. a.i./A) to control annual bluegrass in seedling.

**FOR USE IN ROUNDUP READY FIELD CORN (BURNDOWN)**

GRASS SPECIES	WEED SIZE (Inches)	Rate when applied alone or with glyphosate
Field Corn	Up to 12	3 fl. oz./A (0.047 lb. a.i.)

• For control of existing stand of Roundup Ready field corn or volunteer Roundup Ready field corn prior to replanting corn.  
 • Care must be taken to avoid in-field boom (spray) overlaps or excessive crop injury may occur.  
 • Replant no sooner than 6 days after application.  
 • Adjuvant directions: NIS at 0.25% v/v plus AMS at 2.5 to 4 lbs./A.  
 • **DO NOT** use COC or MSD with **Tigris PS Clethodim** in this use pattern.

**Restrictions:**

- **DO NOT** apply more than 3 fl. oz./A (0.05 lb. a.i./A) per application.
- **DO NOT** apply more than 12 fl. oz./A (0.19 lb. a.i./A) per year.
- **DO NOT** apply more than 4 applications per year.
- For repeat applications, make on a minimum of a 14-day interval.

**DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES IN CANOLA, DRIED SHELLED BEAN & PEA (INCLUDING SOYBEAN), EDIBLE PODDED LEGUME VEGETABLES, FLAX, MUSTARD SEED, SUCCULENT BEAN & PEA, AND SUGAR BEET  
(REDUCED RATE DIRECTIONS NOT FOR USE IN CALIFORNIA.)**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture, excessive moisture, low or high temperatures and/or under very low humidity.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (Inches)	RATE FL. OZ./ACRE <sup>(1)</sup>
Barnyardgrass	<i>Echinochloa crus-galli</i>	1 to 4	4 (0.0625 lb. a.i.)
Broodleaf Signalgrass	<i>Bracharia platyphylla</i>	1 to 4	5 (0.078 lb. a.i.)
Crabgrass			
Large	<i>Digitaria sanguinalis</i>	1 to 3*	4 (0.0625 lb. a.i.)
Large	<i>Digitaria sanguinalis</i>	1 to 4*	5 (0.078 lb. a.i.)
Smooth	<i>Digitaria ischaemum</i>	1 to 3*	4 (0.0625 lb. a.i.)
Smooth	<i>Digitaria ischaemum</i>	1 to 4*	5 (0.078 lb. a.i.)
Southern	<i>Digitaria ciliaris</i>	1 to 4*	5 (0.078 lb. a.i.)
Fall Panicum	<i>Panicum dichotomiflorum</i>	1 to 4	4 (0.0625 lb. a.i.)

\* Length of lateral growth

(1) Always add a crop oil concentrate at 1 qt./A by ground application to the finished spray volume.

(continued)

## DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES IN CANOLA, DRIED SHELLED BEAN &amp; PEA (INCLUDING SOYBEAN), EDIBLE PODDED LEGUME VEGETABLES, FLAX, MUSTARD SEED, SUCCULENT BEAN &amp; PEA, AND SUGAR BEET (cont'd)

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT (inches)	RATE FL. OZ./ACRE <sup>(1)</sup>
Foxtail			
Giant	<i>Setaria faberii</i>	1 to 4	4 (0.0625 lb. a.i.)
Green	<i>Setaria viridis</i>	1 to 4	4 (0.0625 lb. a.i.)
Millet	<i>Setaria italica</i>	1 to 4	5 (0.078 lb. a.i.)
Yellow	<i>Setaria glauca</i>	1 to 4	4 (0.0625 lb. a.i.)
Seedling Johnsongrass	<i>Sorghum halepense</i>	1 to 6	5 (0.078 lb. a.i.)
Shattercane	<i>Sorghum bicolor</i>	4 to 10	4 (0.0625 lb. a.i.)
Texas Panicum	<i>Panicum texanum</i>	1 to 4	5 (0.078 lb. a.i.)
Volunteer Cereals			
Barley	<i>Hordeum vulgare</i>	1 to 4	5 (0.078 lb. a.i.)
Oats	<i>Avena sativa</i>	1 to 4	5 (0.078 lb. a.i.)
Wheat	<i>Triticum aestivum</i>	1 to 4	5 (0.078 lb. a.i.)
Volunteer Corn**	<i>Zea mays</i>	4 to 12	4 (0.0625 lb. a.i.)
Wild Proso Millet	<i>Panicum miliaceum</i>	1 to 6	4 (0.0625 lb. a.i.)
Wild Oats	<i>Avena fatua</i>	1 to 4	5 (0.078 lb. a.i.)

\*\* Not S.R. Corn

(1) Always add a crop oil concentrate at 1 qt./A by ground application to the finished spray volume.

#### DIRECTIONS FOR PERENNIAL GRASSES

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

#### Restrictions:

- DO NOT exceed the maximum rate per application listed in the table **CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM**.
- DO NOT exceed the maximum number of applications per year listed in the table **CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM**.
- DO NOT exceed the maximum yearly rate listed in the table **CROP SPECIFIC DIRECTIONS AND RESTRICTIONS FOR TIGRIS PS CLETHODIM**.
- For repeat applications, make on a minimum of a 14-day interval.

GRASS SPECIES	WEED HEIGHT (inches)	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE
Bermudagrass ( <i>Cynodon dactylon</i> )			
First Application	3 (or up to 6" runners)	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	3 (or up to 6" runners)	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Fescue, tall ( <i>Festuca arundinacea</i> )			
First Application	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Foxtail Barley ( <i>Hordeum jubatum</i> )			
First Application	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Orchardgrass ( <i>Dactylis glomerata</i> )			
First Application	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Quackgrass* ( <i>Elytrigia repens</i> )			
First Application	4 to 12	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	4 to 12	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Rhizome Johnsongrass ( <i>Sorghum halepense</i> )			
First Application	12 to 24	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	6 to 18	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Wirestem Muhly ( <i>Muhlenbergia frondosa</i> )			
First Application	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)

\*Control of quackgrass, perennial bluegrass and bentgrass with **Tigris PS Clethodim** may be enhanced by adding AMS at 2.5 to 4 lbs./A.

(continued)

**DIRECTIONS FOR PERENNIAL GRASSES (cont'd)**

GRASS SPECIES	WEED HEIGHT (inches)	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE
Perennial Bluegrass*			
Boroughstalk ( <i>Poa trivialis</i> )			
Kentucky ( <i>Poa pratensis</i> )			
First Application	2 to 4	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	2 to 4	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
 Bentgrass* ( <i>Agrostis spp.</i> )			
First Application	2 to 4	-	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	2 to 4	-	16 (0.25 lb. a.i.)

\*Control of quackgrass, perennial bluegrass and bentgrass with **Tigris PS Clethodim** may be enhanced by adding AMS at 2.5 to 4 lbs./A.

**TANK MIXES INFORMATION**

The labels for each of the herbicides directed for tank mixing with **Tigris PS Clethodim** are unique to the characteristics of those products and contain restrictions and limitations that may be more restrictive than the **Tigris PS Clethodim** label in certain considerations.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Those concerns may include, but are not limited to:

1. Geographic restrictions – all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
2. Crop rotation restrictions;
3. Applicator certification requirements;
4. Worker safety rules (including protective clothing, reentry time, posting);
5. Soil type or soil characteristics (including pH, OM);
6. Maximum dosage or number of applications per year;
7. Rain free period required; or
8. Application timing (including pre-harvest interval).
9. **DO NOT** exceed the total year rate.

**THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.**

**TANK MIX APPLICATION OF TIGRIS PS CLETHODIM AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS**

- Apply only to actively growing grass and broadleaf weeds at specified height or growth stage listed on each label.
- Apply when the first grass or broadleaf weed species in a mixed population reaches the specified height or growth stage for treatment.
- Apply under favorable soil moisture and humidity that exist a few days after rainfall or within 7 days after irrigation.
- Always add the appropriate adjuvant to the spray mix at the rate specified for each specific tank mix combination.
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs, or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim**, as specified in the respective size and rate tables.
- **DO NOT** tank mix **Tigris PS Clethodim** when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.

**MIXING INSTRUCTIONS**

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, add the correct amount of **Tigris PS Clethodim**. Agitation must create a rippling or rolling action on the water surface.
3. If tank mixing **Tigris PS Clethodim** with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates, and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants (crop oil concentrate, non-ionic surfactant and/or nitrogen solution).
5. Fill spray tank to desired level with water. **Agitation must continue until all spray solution has been applied.**

Failure to agitate the spray solution may result in improper mixing of the herbicides and unsatisfactory weed control. Mixing and compatibility qualities must be verified by a jar test.

**INFORMATION ON ANTAGONISM**

Tank mixes of **Tigris PS Clethodim** with post-emergence broadleaf herbicides have shown some reduction or failure to control certain grass species which would have otherwise been controlled when **Tigris PS Clethodim** is applied alone. Activity of the post-emergence broadleaf herbicide in the tank mix is not affected.

**DIRECTIONS FOR ALFALFA**

**Table 1. TIGRIS PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR ALFALFA**

(Refer to tables above for specific grasses and growth stages.)

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
<b>Tigris PS Clethodim</b> + Dimethylamine 4-(2,4-dichlorophenoxy) butyrate <sup>(4)</sup>	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) + Refer to Dimethylamine 4-(2,4-dichlorophenoxy) butyrate label for use rate.	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) + Refer to Dimethylamine 4-(2,4-dichlorophenoxy) butyrate label for use rate.	1%	1%
<b>Tigris PS Clethodim</b> + imazethapyr <sup>(5)</sup>	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) + Refer to imazethapyr labels for use rate.	-	1%	1%
<b>Tigris PS Clethodim</b> + bromoxynil <sup>(6)(7)</sup>	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) + Refer to bromoxynil label for use rate.	-	0.5%	0.5%

- (1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.
- (2) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.
- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- (4) **Tigris PS Clethodim** plus Dimethylamine 4-(2,4-dichlorophenoxy) butyrate may increase the severity of crop injury when tank mixed. Alfalfa plants will generally outgrow this temporary crop injury within a few weeks.
- (5) Before using this tank mix, read and understand the imazethapyr labels for geographical restrictions and restrictions regarding alfalfa growth stage and type. Failure to do so can result in crop injury to alfalfa. **DO NOT** feed, graze, or harvest alfalfa for 30 days following an application of imazethapyr to alfalfa.
- (6) In the states of Washington, Oregon, Idaho, Montana, Wyoming, Colorado, Utah, Nevada and the western halves of North Dakota, South Dakota, Nebraska, and Kansas: The **Tigris PS Clethodim** plus bromoxynil tank mix must be applied in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 2 trifoliates. Unacceptable crop injury may occur to alfalfa seedlings less than the 2 trifoliolate leaf stage. **Tigris PS Clethodim** plus bromoxynil applications made when temperatures are expected to exceed 80°F at (and 3 days following) application can result in unacceptable crop injury. In the states not listed above, apply in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 4 trifoliolate leaves. When the alfalfa stand is uneven and conditions favor leaf burn, unacceptable crop injury may occur to alfalfa in the 2 trifoliolate or smaller stage of growth. **Tigris PS Clethodim** plus bromoxynil applications made when temperatures are expected to exceed 70°F at (and 3 days following) application can result in unacceptable crop injury. Crop leaf burn can occur following **Tigris PS Clethodim** plus bromoxynil application. Warm, humid conditions may enhance leaf burn. New crop growth will not be affected.
- (7) **DO NOT** apply when alfalfa is under moisture, temperature, insect, or disease stress or has been stressed by other pesticide carryover or application.

**DIRECTIONS FOR CANOLA (EXCEPT FLAX)**

[Rapeseed Subgroup 20A (except flax seed, mustard seed, and sesame seed)]

**Table 2. REDUCED RATE TIGRIS PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR CANOLA**

(Refer to the tables above for specific grasses and growth stages.)

PRODUCT	APPLICATION RATES/ACRE		AMMONIUM SULFATE	
	ANNUAL GRASSES <sup>(1)</sup>	PERENNIAL GRASSES	GROUND	AIR
<b>Tigris PS Clethodim<sup>(2)</sup></b> + glufosinate <sup>(3)</sup>	4 to 5 fl. oz. (0.063-0.078 lb. a.i.) + Refer to glufosinate label for use rate.	-	3.0 lbs.	3.0 lbs.

(1) Annual grasses and sizes controlled with these tank mixtures are those that are identified in the **DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES** table.

(2) **DO NOT** apply **Tigris PS Clethodim** tank mix during or after bolting or flowering or crop injury may occur.

(3) For use only on LibertyLink® canola.

DIRECTIONS FOR COTTON

Table 3. TIGRIS PS CLETHODIM TANK MIXED WITH LACTOFEN AND MSMA APPLIED POST DIRECTED TO COTTON

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> V/V	COMMENTS
	ANNUAL GRASSES	PERENNIAL GRASSES		
Tigris PS Clethodim <sup>(4)</sup> + lactofen + MSMA (4 lbs./gal.) or MSMA (6.6 lbs./gal.)	6 to 8 fl. oz. (0.094-0.125 lb. a.i.)	8 to 16 fl. oz. (0.125-0.25 lb. a.i.)	1%	Reduce broadcast rate in proportion to the band area actually treated.
	See lactofen label for rates to control broadleaf weeds and height limitations for cotton. Refer to the <b>Tigris PS Clethodim</b> label for weed height and species controlled.			
	See MSMA label for rates to control broadleaf weeds and height limitations for cotton. Refer to the <b>Tigris PS Clethodim</b> label for weed height and species controlled.			

- (1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.
- (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.
- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.
- (4) If at the time of application, grass height is so tall that post-directed applications cannot get good coverage over the top of the grassy weeds, then poor control may result and a second (non-post directed) application of **Tigris PS Clethodim** may be necessary.

Table 4. TIGRIS PS CLETHODIM TANK MIXED WITH BROMOXYNIL TO CONTROL EMERGED WEEDS IN BXM COTTON AS A BROADCAST APPLICATION

PRODUCT <sup>(1)</sup>	APPLICATION RATES/ACRE <sup>(2)</sup>		CROP OIL CONCENTRATE PER ACRE <sup>(3)</sup>	COMMENTS <sup>(7)</sup>
	ANNUAL GRASSES	PERENNIAL GRASSES		
Tigris PS Clethodim + bromoxynil <sup>(4,5,6)</sup>	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + See bromoxynil label for rates to control broadleaf weeds and height limitations for cotton.		1 qt.	See charts for grasses controlled.

- (1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage.
- (2) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** at the specified rate with the appropriate amount of crop oil concentrate in a non-bromoxynil tank mix.
- (3) Always add a crop oil concentrate at 1 qt./A by ground in the finished spray solution.
- (4) Applications of bromoxynil can be made only to cotton that has been genetically modified for crop resistance to post-emergence over-the-top application of bromoxynil.
- (5) **DO NOT** apply the **Tigris PS Clethodim** plus bromoxynil tank mix within 75 days of harvest.
- (6) **DO NOT** exceed 2 applications of bromoxynil before cotton is 12 inches tall and 1 application after 12 inches tall.
- (7) Use a minimum of 10 gals. of spray solution per acre.

Table 5. TIGRIS PS CLETHODIM TANK MIXED WITH GLYPHOSATE TO CONTROL EMERGED GRASSES IN COTTON AS A BROADCAST APPLICATION

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		ADJUVANT		COMMENTS
	ANNUAL GRASSES	PERENNIAL GRASSES	Glyphosate formulation with built in adjuvant	Glyphosate formulation without built in adjuvant	
Tigris PS Clethodim + glyphosate	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) See glyphosate label for rates to control broadleaf weeds and height limitations for cotton.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.)	Non-ionic surfactant @ 0.125 to 0.25% v/v plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier	Crop oil concentrate @ 1 pt./A plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals. of carrier	See charts for grasses controlled. Use a minimum of 10 gals. of spray solution per acre.

- (1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** at the specified rate with the appropriate amount of crop oil concentrate.

**DIRECTIONS FOR DRY SHELLED AND SUCCULENT BEANS**  
**[Succulent Shelled Pea and Beans Subgroup 6B]**  
**[Dried Shelled Pea and Bean (except Soybean 6C)]**

**Table 6. TIGRIS PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR DRY SHELLED AND SUCCULENT BEANS**  
 (Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
Tigris PS Clethodim + bentazon	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to bentazon label for use rate.	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) + Refer to bentazon label for use rate.	1%	1%

- (1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.
- (2) Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.
- (3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

**DIRECTIONS FOR FLAX**

**Table 7. REDUCED RATE TIGRIS PS CLETHODIM TANK MIXED WITH BROADLEAF HERBICIDES FOR FLAX**  
 (Refer to the tables above for specific grasses and growth stages.)

PRODUCT	APPLICATION RATES/ACRE		CROP OIL CONCENTRATE	
	ANNUAL GRASSES <sup>(1)</sup>	PERENNIAL GRASSES	GROUND	AIR
Tigris PS Clethodim + MCPA and bromoxynil octanoate and bromoxynil heptanoate	4 to 5 fl. oz. (0.063-0.078 lb. a.i.) + Refer to product label for use rate.	-	-	1 pt. 1 pt.
Tigris PS Clethodim + bromoxynil <sup>(2,3)</sup>	4 to 5 fl. oz. (0.063-0.078 lb. a.i.) + Refer to bromoxynil label for use rate.	-	-	1 pt. 1 pt.
Tigris PS Clethodim + MCPA ester <sup>(2,3)</sup>	4 to 5 fl. oz. (0.063-0.078 lb. a.i.) + Refer to MCPA ester label for use rate.	-	-	1 pt. 1 pt.

- (1) Annual grasses and sizes controlled with these tank mixtures are those that are identified in the **DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES** table.
- (2) **DO NOT** apply **Tigris PS Clethodim** tank mix during or after the bud stage or to ornamental flax or crop injury may occur.
- (3) **DO NOT** apply tank mixes if temperatures are expected to exceed 85°F at (or 3 days following) application or crop injury may occur.

**DIRECTIONS FOR SOYBEAN**

**Table 8. TIGRIS PS CLETHODIM TANK MIXES<sup>(3)</sup> TO CONTROL ANNUAL GRASSES WHEN USED AS A BURNDOWN IN NO-TILL SOYBEAN**

PRODUCT	PRODUCT RATE/ACRE <sup>(1)</sup>	GRASS HEIGHT (inches)	CROP OIL CONCENTRATE /ACRE <sup>(2)</sup>	28%N OR 32%N QTS./A OR 2.5 TO 4.0 LBS. AMS
Tigris PS Clethodim + 2,4-D ester <sup>(3)</sup>	3 fl. oz. (0.047 lb. a.i.)	Foxtail 1 to 3 Fall Panicum 1 to 3	1 qt.	1 to 2 qts. or 2.5 to 4 lbs. AMS
	4 fl. oz. (0.063 lb. a.i.)	Foxtail 1 to 4 Fall Panicum 1 to 4	1 qt.	1 to 2 qts. or 2.5 to 4 lbs. AMS
	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to 2,4-D ester label for use rate.	(See Grass Chart for grasses claimed.)	1 qt.	1 to 2 qts. or 2.5 to 4 lbs. AMS

\* 2,4-D ester must not be used where drift sensitive crops may be grown.

- (1) If regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** according to the appropriate size and rate directions.
- (2) Always use a crop oil concentrate at the listed rate in the finished spray volume.
- (3) The following products can be tank mixed with **Tigris PS Clethodim** plus 2,4-D ester: flumioxazin, chlorimuron-ethyl + sulfentrazone, metolachlor, s-metolachlor, pendimethalin, metribuzin, metribuzin plus the metolachlor/s-metolachlor products, and glyphosate.

Table 9. TIGRIS PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN

(Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(2)</sup>	ANNUAL GRASSES	APPLICATION RATES/ACRE <sup>(1)</sup> PERENNIAL GRASSES	CROP OIL CONCENTRATE <sup>(3)</sup> (V/V) GROUND	AIR
Tigris PS Clethodim + lactofen	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the lactofen label for use rate.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + Refer to the lactofen label for use rate.	0.5 to 1%	1%
Tigris PS Clethodim + bentazon	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to the bentazon label for use rate.	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) + Refer to the bentazon label for use rate.	1%	1%
Tigris PS Clethodim + glyphosate (For use on Roundup Ready soybean only)	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the glyphosate label for use rate.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + Refer to the glyphosate label for use rate.	0.5 to 1% <sup>(4)</sup>	1% <sup>(4)</sup>
Tigris PS Clethodim + acifluorfen	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the acifluorfen label for use rate.	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the acifluorfen label for use rate.	0.5% to 1%	1%
Tigris PS Clethodim + fomesafen <sup>(6)</sup>	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the fomesafen label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + Refer to the fomesafen label for use rates.	1%	1%
Tigris PS Clethodim + chlormuron-ethyl	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to the chlormuron-ethyl label for use rates.	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) + Refer to the chlormuron-ethyl label for use rates.	1%	1%
Tigris PS Clethodim <sup>(5)</sup> + imazethapyr	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the imazethapyr label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + Refer to the imazethapyr label for use rates.	1%	1%
Tigris PS Clethodim <sup>(5)</sup> + lactofen + chlormuron-ethyl	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to the lactofen label for use rates. + Refer to the chlormuron-ethyl label for use rates.	-	0.5%	1%
Tigris PS Clethodim <sup>(5)</sup> + lactofen + bentazon	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to the lactofen label for use rates. + Refer to the bentazon label for use rates.	-	0.5%	1%

(1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.

(2) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

(4) The addition of 2.5 lbs./A of ammonium sulfate is required when **Tigris PS Clethodim** is tank mixed with glyphosate. If the glyphosate formulation has a stand-alone built in adjuvant, add 0.125% v/v non-ionic surfactant in place of crop oil concentrate. If the glyphosate formulation does not have a built in adjuvant, add 0.5% to 1% crop oil concentrate for ground application and 1% v/v for aerial application.(5) The addition of 1 to 2 pts. of a liquid fertilizer (10-34-0, 28%N, or 32%N) is advised when **Tigris PS Clethodim** is tank mixed with imazethapyr, flumiclorac, bentazon and acifluorfen, cloransulam-methyl, thifensulfuron and chlormuron-ethyl, imazomox, cloransulam-methyl and flumetsulam, lactofen plus chlormuron-ethyl, lactofen plus bentazon, lactofen plus imazethapyr, lactofen plus cloransulam-methyl, lactofen plus thifensulfuron and chlormuron-ethyl, and lactofen plus imazomox. An equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate.

(6) Refer to the fomesafen label for geographic and rotational restrictions.

(continued)

Table 9. TIGRIS PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN (cont'd)

(Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
Tigris PS Clethodim <sup>(5)</sup> + lactofen + imazethapyr	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to the lactofen label for use rates. + Refer to the imazethapyr label for use rates.	-	0.5%	1%
Tigris PS Clethodim <sup>(5)</sup> + flumiclorac + imazethapyr	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to the flumiclorac label for use rates. + Refer to the imazethapyr label for use rates.	-	1%	1%
Tigris PS Clethodim <sup>(5)</sup> + flumiclorac + bentazon	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to the flumiclorac label for use rates. + Refer to the bentazon label for use rates.	-	1%	1%
Tigris PS Clethodim <sup>(5)</sup> + flumiclorac + chlorimuron-ethyl	8 to 10 fl. oz. (0.125-0.156 lb. a.i.) + Refer to the flumiclorac label for use rates. + Refer to the chlorimuron-ethyl label for use rates.	-	1%	1%
Tigris PS Clethodim <sup>(5)</sup> + lactofen + flumiclorac	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the lactofen label for use rates. + Refer to the flumiclorac label for use rates.	-	0.5%	1%
Tigris PS Clethodim <sup>(5)</sup> + cloransulam-methyl	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the cloransulam-methyl label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + Refer to the cloransulam-methyl label for use rates.	1%	-
Tigris PS Clethodim <sup>(5)</sup> + lactofen + cloransulam-methyl	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the lactofen label for use rates. + Refer to the cloransulam-methyl label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + Refer to the lactofen label for use rates. + Refer to the cloransulam-methyl label for use rates.	1%	-
Tigris PS Clethodim <sup>(5)</sup> + imazamox	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the imazamox label for use rates.	-	1%	-

(1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.

(2) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

(4) The addition of 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N) is advised when **Tigris PS Clethodim** is tank mixed with imazethapyr, flumiclorac, bentazon and acifluorfen, cloransulam-methyl, thifensulfuron and chlorimuron-ethyl, imazamox, cloransulam-methyl and flumetsulam, lactofen plus chlorimuron-ethyl, lactofen plus bentazon, lactofen plus imazethapyr, lactofen plus cloransulam-methyl, lactofen plus thifensulfuron and chlorimuron-ethyl, and lactofen plus imazamox. An equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate.

(continued)

Table 9. TIGRIS PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN (cont'd)

(Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	AIR
	ANNUAL GRASSES	PERENNIAL GRASSES		
Tigris PS Clethodim <sup>(5)</sup> + lactofen + imazamox	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the lactofen label for use rates. + Refer to the imazamox label for use rates.		1%	-
Tigris PS Clethodim <sup>(5)</sup> + chlorimuron-ethyl and thifensulfuron	6 to 8 fl. oz. <sup>(7)</sup> (0.094-0.125 lb. a.i.) + Refer to the chlorimuron-ethyl and thifensulfuron label for use rates.		1 qt.	-
Tigris PS Clethodim <sup>(5)</sup> + lactofen + chlorimuron-ethyl and thifensulfuron	6 to 8 fl. oz. <sup>(7)</sup> (0.094-0.125 lb. a.i.) + Refer to the lactofen label for use rates. + Refer to the chlorimuron-ethyl and thifensulfuron label for use rates.		1 pt.	-
Tigris PS Clethodim <sup>(5)</sup> + flumiclorac	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the flumiclorac label for use rates.		1 qt.	-
Tigris PS Clethodim + cloransulam-methyl + fomesafen <sup>(6)</sup>	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the cloransulam-methyl label for use rates. + Refer to the fomesafen label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + Refer to the cloransulam-methyl label for use rates. + Refer to the fomesafen label for use rates.	1%	-

(1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.

(2) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

(5) The addition of 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N) is advised when **Tigris PS Clethodim** tank mix with imazethapyr, flumiclorac, bentazon and acifluorfen, cloransulam-methyl, thifensulfuron and chlorimuron-ethyl, imazamox, cloransulam-methyl and flumetsulam, lactofen plus chlorimuron-ethyl, lactofen plus bentazon, lactofen plus imazethapyr, lactofen plus cloransulam-methyl, lactofen plus thifensulfuron and chlorimuron-ethyl, and lactofen plus imazamox. An equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate.

(6) Refer to the fomesafen label for geographic and rotational restrictions.

(7) Annual grasses and sizes controlled with these tank mixes are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table.

Table 10. REDUCED RATE TIGRIS PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN (Refer to table for reduced rate use in canola, dry shelled bean &amp; pea, edible podded legume vegetables, flax, mustard seed, soybean, succulent bean &amp; pea and sugar beet for small annual grasses for specific grasses and growth stages.)

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3,4)</sup> (V/V)	
	ANNUAL GRASSES <sup>(2)</sup>	PERENNIAL GRASSES	GROUND	AIR
Tigris PS Clethodim + cloransulam-methyl	4 to 8 fl. oz. (0.063-0.125 lb. a.i.) + Refer to the cloransulam-methyl label for use rates.	-	1%	1%
Tigris PS Clethodim + imazethapyr	4 to 6 fl. oz. (0.063-0.094 lb. a.i.) + Refer to the imazethapyr label for use rates.	-	1%	1%

(1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.

(2) Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE TO CONTROL SMALL ANNUAL GRASSES table.

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

(4) The addition of 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N, or 32%N) is required when **Tigris PS Clethodim** tank mix with reduced rates. An equivalent amount (2.5 to 4 lbs./A) of spray grade ammonium sulfate (AMS) may be added in place of liquid fertilizer. Fertilizer adjuvants are to be added in addition to the crop oil concentrate.

**DIRECTIONS FOR PEANUT (INCLUDING PERENNIAL)**

**Table 11. TIGRIS PS CLETHODIM TANK MIXES WITH BROADLEAF HERBICIDES FOR PEANUT (INCLUDING PERENNIAL)**

(Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES <sup>(2)</sup>	PERENNIAL GRASSES	GROUND	AIR
Tigris PS Clethodim + bentazon	8 to 10 fl. oz. (0.125–0.156 lb. a.i.) + Refer to the bentazon label for use rates.	-	1%	1%
Tigris PS Clethodim + acifluorfen	8 to 10 fl. oz. (0.125–0.156 lb. a.i.) + Refer to the acifluorfen label for use rates.	-	1%	1%

(1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.

(2) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

**GRASS SUPPRESSION FOR HARVEST EFFICIENCY IN PEANUT (INCLUDING PERENNIAL) WITH TIGRIS PS CLETHODIM**

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ACRE	HIGH RATE FL. OZ./ACRE
Annual and perennial grasses that exceed height claimed for control on height charts "ANNUAL GRASSES" and "PERENNIAL GRASSES"	Up to and including grasses in the seed head stage.	16 (0.25 lb. a.i.)	32 (0.5 lb. a.i.)

**DO NOT** apply as part of a tank mix when applying **Tigris PS Clethodim** for grass suppression.

Add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.

**DIRECTIONS FOR SUGAR BEET**

**Table 12. TIGRIS PS CLETHODIM TANK MIXED WITH CLOPYRALID APPLIED TO SUGAR BEET**

(Refer to the tables above for specific grasses and growth stages.)

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)	
	ANNUAL GRASSES	PERENNIAL GRASSES	GROUND	AIR
Tigris PS Clethodim + clopyralid	6 to 8 fl. oz. (0.094–0.125 lb. a.i.) See the clopyralid label for rates.	8 to 16 fl. oz. (0.125–0.25 lb. a.i.)	1%	1%

(1) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.

(2) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

**Table 13. TIGRIS PS CLETHODIM TANK MIXED WITH DESMEDIPHAM + PHENMEDIPHAM OR DESMEDIPHAM APPLIED TO SUGAR BEET**

PRODUCT <sup>(1)</sup>	WEEDS CONTROLLED		WEED HEIGHT (Inches)	APPLICATION RATE/ACRE <sup>(2)</sup>
	COMMON NAME	SCIENTIFIC NAME		
Tigris PS Clethodim <sup>(3)</sup> + desmedipham + phenmediphham or desmedipham	Barnyardgrass Foxtail Foxtail Millet Wild Oat Wild Proso Millet	<i>Echinochloa crus-galli</i> <i>Setaria spp.</i> <i>Setaria italica</i> <i>Avena fatua</i> <i>Panicum miliaceum</i>	1 to 3 1 to 3 1 to 3 1 to 3 1 to 3	8 fl. oz. (0.125 lb. a.i.)
				See desmedipham + phenmediphham label for rates to control broadleaf weeds. <b>No additives are advised in the tank mix.</b>
				See desmedipham label for rates to control broadleaf weeds. <b>No additives are advised in the tank mix.</b>

(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.

(2) **DO NOT** use crop oil concentrate. No additives are advised in the tank mix. If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix herbicide), according to the appropriate size and rate directions.

(3) If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** at full label rate with appropriate rate of crop oil concentrate.

Table 14. TIGRIS PS CLETHODIM PLUS DESMEDIPHAM • PHENMEDIPHAM OR DESMEDIPHAM TANK MIX FOR 3 SEQUENTIAL APPLICATIONS FOR ANNUAL GRASS CONTROL (MICRO-RATE APPLICATION) IN SUGAR BEET

PRODUCT	APPLICATION RATES/ACRE <sup>(1)</sup>		METHYLATED SEED OIL <sup>(2)</sup> (V/V)	
	ANNUAL GRASSES	GRASSES CONTROLLED (Inches)	GROUND	AIR
Tigris PS Clethodim + desmedipham + phenmedipham or desmedipham	2 to 3 fl. oz. (0.031-0.047 lb. a.i./A) + Refer to product labels for use rates.	Green Foxtail (1-2) Yellow Foxtail (1-2) Barnyardgrass (1-2) Wild Oat (1-2) Volunteer Cereals (1-2)	1.5%	1.5%

(1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not advised in these situations.

(2) Always use a methylated seed oil at the listed rate (but not less than 1 pt./A) in the finished spray volume.

#### Directions for Use for Micro-Rate Applications to Sugar Beet

##### Application Information

Multiple micro-rate applications of **Tigris PS Clethodim** in tank mixtures with reduced rates of desmedipham + phenmedipham or desmedipham and methylated seed oils may be applied by air or ground equipment to sugar beet to control early germinating annual grasses listed above. The rate of desmedipham + phenmedipham or desmedipham must not exceed 0.12 lb. a.i./A (broadcast application) when in combination with these spray adjuvants. Note that maximum rate allowed varies depending on crop growth stage. The use of wetting agents or spray adjuvants with conventional rates (0.75 to 1.25 lbs. a.i./A) or multiple low rate (0.24 to 0.75 lb. a.i./A) applications of desmedipham + phenmedipham or desmedipham is prohibited on the desmedipham + phenmedipham or desmedipham master label. Favorable climatic conditions (good conditions for plant growth and development) are essential for adequate weed control. All use precautions and restrictions on the desmedipham + phenmedipham or desmedipham master labels must be followed.

#### Directions for Using Micro-Rate Multiple Applications of Tigris PS Clethodim Tank Mixes

Apply **Tigris PS Clethodim** in broadcast applications only at a rate of 2 to 3 fl. oz./A (0.031-0.047 lb. a.i./A) in tank mixture with either desmedipham + phenmedipham or desmedipham following the directions for use on the tank mix partner label. A minimum of 3 sequential applications of 2 fl. oz./A (0.031 lb. a.i./A) or a minimum of 2 sequential applications of 3 fl. oz./A (0.047 lb. a.i./A) must be utilized for **Tigris PS Clethodim** tank mixtures. A minimum of 3 sequential applications of desmedipham + phenmedipham or desmedipham must be used. Accurate timing is essential; make initial application immediately after weeds emerge, and make repeat applications on 5- to 7-day intervals. If weed control is not adequate due to climatic conditions, spray coverage or other factors, return to conventional application rates of **Tigris PS Clethodim** (6 to 8 fl. oz./A (0.094-0.125 lb. a.i.)) and add rates of desmedipham + phenmedipham or desmedipham as directed on their label. When using conventional rates of desmedipham + phenmedipham or desmedipham in tank mixtures with **Tigris PS Clethodim**, a spray adjuvant is not advised.

#### Use Precautions for Micro-Rate Applications: (See **Tigris PS Clethodim**, desmedipham + phenmedipham and desmedipham master label for further use precautions.)

- Not all weeds will be adequately controlled, even with favorable climatic conditions. Conventional rates of **Tigris PS Clethodim**, desmedipham + phenmedipham or desmedipham and/or hand labor may be required if multiple micro-rate applications do not adequately control weeds. Plugging of spray nozzles may be encountered due to the potential for formation of a precipitate in the spray solution that is often associated with micro-rate applications. To the extent consistent with applicable law, Tigris, LLC will not be responsible for any nozzle plugging that may occur with the use of multiple micro-rate applications. Methylated seed oils must not be added if the desmedipham + phenmedipham or desmedipham rate exceeds 0.12 lb. a.i./A broadcast, as the addition of methylated seed oils could increase the possibility of crop injury at dosage rates greater than 0.12 lb. a.i./A.
- Ground Application:** Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 10 gals. and a maximum of 20 gals. of spray solution per acre. Spray pressures must reflect a minimum of 30 PSI and a maximum of 60 PSI at the nozzle. **DO NOT** use flood nozzles.
- Aerial Application:** Use of sufficient spray volumes is essential to ensure complete coverage. Use a minimum of 5 gals. and a maximum of 15 gals. of spray solution per acre.

Table 15. TANK MIX APPLICATION OF TIGRIS PS CLETHODIM AND FUNGICIDES FOR CONTROL OF GRASS WEEDS AND DISEASES IN SUGAR BEET

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE <sup>(3)</sup> (V/V)
	ANNUAL GRASSES	PERENNIAL GRASSES	
Tigris PS Clethodim + tetraconazole	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) + Refer to the tetraconazole label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) + Refer to the tetraconazole label for use rates.	1%

(1) If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix fungicide), according to the appropriate size and rate directions.

(2) Refer to **Tigris PS Clethodim** and fungicide label for rates and weeds and diseases controlled.

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

Table 16. TANK MIX APPLICATION OF TIGRIS PS CLETHODIM AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA, COTTON, MINT (PEPPERMINT AND SPEARMINT TOPS), PEANUT (INCLUDING PERENNIAL), SOYBEAN AND SUNFLOWER

PRODUCT <sup>(2)</sup>	APPLICATION RATES/ACRE <sup>(1)</sup>		CROP OIL CONCENTRATE (V/V) <sup>(3)</sup>	CROP					
	ANNUAL GRASSES	PERENNIAL GRASSES		Mint <sup>(4)</sup>	Cotton	Mint <sup>(4,5)</sup>	Peanut	Soybean	Sunflower
Tigris PS Clethodim + acephate	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) * Refer to the acephate label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) * Refer to the acephate label for use rates.	1%		X	X	X	X(6)	
Tigris PS Clethodim + fenpropathrin	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) * Refer to the fenpropathrin label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) * Refer to the fenpropathrin label for use rates.	1%		X		X		
Tigris PS Clethodim + esfenvalerate	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) * Refer to the esfenvalerate label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) * Refer to the esfenvalerate label for use rates.	1%						X
Tigris PS Clethodim + lambda-cyhalothrin	6 to 8 fl. oz. (0.094-0.125 lb. a.i.) * Refer to the lambda-cyhalothrin label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.) * Refer to the lambda-cyhalothrin label for use rates.	1%						X
Tigris PS Clethodim + lambda-cyhalothrin	10 to 16 fl. oz. <sup>(7)</sup> (0.156-0.25 lb. a.i.) * Refer to the lambda-cyhalothrin label for use rates.	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) * Refer to the lambda-cyhalothrin label for use rates.	1%		X				
Tigris PS Clethodim + B-cyfluthrin	10 to 16 fl. oz. <sup>(7)</sup> (0.156-0.25 lb. a.i.) * Refer to the B-cyfluthrin label for use rates.	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) * Refer to the B-cyfluthrin label for use rates.	1%		X				
Tigris PS Clethodim + dimethoate	10 to 16 fl. oz. <sup>(7)</sup> (0.156-0.25 lb. a.i.) * Refer to the dimethoate label for use rates.	10 to 16 fl. oz. <sup>(7)</sup> (0.156-0.25 lb. a.i.) * Refer to the dimethoate label for use rates.	1%		X				
Tigris PS Clethodim + chlorpyrifos	10 to 16 fl. oz. <sup>(7)</sup> (0.156-0.25 lb. a.i.) * Refer to the chlorpyrifos label for use rates.	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) * Refer to the chlorpyrifos label for use rates.	1 to 2 pts. <sup>(8)</sup>		X				
Tigris PS Clethodim + permethrin	10 to 16 fl. oz. <sup>(7)</sup> (0.156-0.25 lb. a.i.) * Refer to the permethrin label for use rates.	10 to 16 fl. oz. (0.156-0.25 lb. a.i.) * Refer to the permethrin label for use rates.	1%		X				

\* Peppermint and Spearmint tops

(1) If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim** alone (without a tank mix fungicide), according to the appropriate size and rate directions.

(2) Refer to **Tigris PS Clethodim** and fungicide label for rates and weeds and diseases controlled.

(3) Always use a crop oil concentrate at the listed rate (but not less than 1 pt./A) in the finished spray volume.

(4) Certain insecticides may cause temporary phytotoxic symptoms on alfalfa and mint (peppermint and spearmint tops) foliage. Refer to the insecticide label for further information. It is suggested that prior to using any of these insecticides/herbicide tank mixtures, that a small area of the field be treated first and observations for crop injury be made prior to treating the whole field.

(5) The **Tigris PS Clethodim** rate must be 6 to 8 fl. oz./A (0.094-0.125 lb. a.i./A) for annual grass control in baby mint (peppermint and spearmint tops), minimum of 8 fl. oz./A (0.125 lb. a.i./A) for annual grass control in established mint (peppermint and spearmint tops) and 8 to 16 fl. oz./A (0.094-0.125 lb. a.i./A) for perennial grass control. Crop oil concentrate must be added at the rate of 1 to 2 pts./A.

(6) Insecticide tank mix use with acephate in soybean is permitted only in a State having an approved Section 2(a) registration for acephate use in soybean.

(7) The **Tigris PS Clethodim** rate must be 6 to 8 fl. oz./A (0.094-0.125 lb. a.i./A) for annual grass control in seedling alfalfa.

(8) For the **Tigris PS Clethodim** plus chlorpyrifos tank mix, reduce the adjuvant rate down to 1 pt./A when the chlorpyrifos rate is 1 pt./A or higher.

Table 17. ROUNDUP READY VOLUNTEER CORN CONTROL IN ROUNDUP READY SOYBEAN WITH TIGRIS PS CLETHODIM TANK MIX

Roundup Ready Volunteer Corn Height (Inches)	Tigris PS Clethodim Rate fl. oz./A	Glyphosate <sup>(1)</sup> rate for formulations with built in adjuvant	Adjuvant
42	4 (0.063 lb. a.i.)	Refer to product label for use rates.	Non-ionic surfactant @ 0.125 to 0.25% v/v plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals./of carrier
12 to 18	5 (0.078 lb. a.i.)		
18 to 24	6 (0.094 lb. a.i.)		

Roundup Ready Volunteer Corn Height (Inches)	Tigris PS Clethodim Rate fl. oz./A	Glyphosate <sup>(1)</sup> rate for formulations without built in adjuvant	Adjuvant
42	4 (0.063 lb. a.i.)	Refer to product label for use rates.	Crop oil concentrate @ 1 pt./A plus ammonium sulfate @ 8.5 to 17 lbs. per 100 gals./of carrier
12 to 18	5 (0.078 lb. a.i.)		
18 to 24	6 (0.094 lb. a.i.)		

(1) Glyphosate formulation must be labeled for use on Roundup Ready soybean.

**THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.**

- Apply only to actively growing grass and broadleaf weeds at specified height or growth stage listed on each label.
- Apply under favorable soil moisture and humidity which exist a few days after rainfall or within 7 days after irrigation.
- Tank mix applications may sometimes result in reduced grass control. If regrowth occurs, or an additional flush of new grass emerges, make a second application of **Tigris PS Clethodim**, as specified in the respective size and rate tables.
- This tank mix may be applied post-emergence to ROUNDUP READY soybean up through the full flowering stage. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants and trees, other than soybean with the ROUNDUP READY gene as severe injury or destruction will result.

**Restrictions:**

- **DO NOT** tank mix **Tigris PS Clethodim** when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.
- **DO NOT** apply less than 60 days before harvest.
- **DO NOT** allow the **Tigris PS Clethodim** plus ROUNDUP to mist, drip, drift or splash onto desirable vegetation as minute quantities of the tank mix can cause severe damage or destruction to the crops, plants or other areas on which treatment was not intended. The likelihood of injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions that allow spray drift to occur including combinations of spray pressure and nozzle type that will result in fine particles (mist) that are likely to drift.

**DIRECTIONS FOR FALLOW LAND**

**Tigris PS Clethodim** may be used to control annual and perennial grasses in land that has been left fallow the previous year and other non-producing agricultural areas. Apply **Tigris PS Clethodim** at 6 to 8 fl. oz./A (0.094-0.125 lb. a.i./A) for annual grasses and 8 to 16 fl. oz./A (0.125-0.25 lb. a.i./A) for perennial grasses. When both grass and broadleaf weeds are the target pest, **Tigris PS Clethodim** may be tank mixed with 2,4-Ester for broad spectrum control. When both annual and perennial grasses occur in the same field, use a minimum of 8 fl. oz./A (0.125 lb. a.i./A) **Tigris PS Clethodim** rate.

**Product Information:**

- Use a minimum spray volume of 5 gals./A for aerial applications and 15 gals./A for ground applications.
- Apply only to actively growing grasses when the first grass reaches the specified weed height as specified by the **ANNUAL** and **PERENNIAL GRASSES** sections of this label.
- Annual grasses that emerge after the **Tigris PS Clethodim** application will not be controlled, and a second application may be necessary.
- The control of perennial grasses may require more than 1 application in non-tilled areas.

**Restrictions:**

- **DO NOT** apply more than 8 fl. oz./A (0.125 lb. a.i./A) per application to Annual Grasses and 16 fl. oz./A (0.25 lb. a.i./A) per application to Perennial Grasses.
- **DO NOT** exceed 5 applications at the 8 fl. oz. (0.094 lb. a.i./A) use rate for Annual Grasses and **DO NOT** exceed 4 applications at the 8 fl. oz. (0.125 lb. a.i./A) rate for Perennial Grasses.
- For repeat applications, make on a minimum of a 14-day interval.
- **DO NOT** apply more than 32 fl. oz./A (0.5 lb. a.i./A) per year.
- **DO NOT** plant any crop for 50 days after application unless clethodim is registered for use in that crop.
- **DO NOT** apply to grasses that have tillered, formed seedheads or exceeded specified growth stage.
- **DO NOT** use flood jet nozzles.
- **DO NOT** apply to drought stressed grasses.
- **DO NOT** mow area for 2 weeks prior to or after the **Tigris PS Clethodim** application.

Table 18. TIGRIS PS CLETHODIM IN TANK MIXES TO CONTROL ANNUAL AND PERENNIAL GRASSES IN FALLOW LAND

PRODUCT	APPLICATION RATES/ACRE(1)		CROP OIL CONCENTRATE(2) (V/V)	GROUND	AIR
	ANNUAL GRASSES	PERENNIAL GRASSES			
Tigris PS Clethodim + 2,4-D ester or dicamba	6 to 8 fl. oz. (0.094-0.125 lb. a.i.)  + Refer to the 2,4-D label for use rates. or See dicamba label for use rates.	8 to 16 fl. oz. (0.125-0.25 lb. a.i.)		1%	1%

(1) Refer to **Tigris PS Clethodim** label for weed height and species control. Review 2,4-D labels for crop restrictions, use rates and weeds controlled.

(2) Always use a crop oil concentrate or methylated seed oil containing at least 15% emulsifier at the listed rate (but not less than 1 pt./A) in the finished spray volume.

Table 19. TIGRIS PS CLETHODIM FOR THE CONTROL AND/OR SUPPRESSION OF TALL FESCUE IN NATIVE PRAIRIE WARM-SEASON GRASS RESTORATION PROJECTS

PRODUCT	PRODUCT RATES	GRASS WEEDS		WEED STAGES
		CONTROLLED/SUPPRESSED	COMMON NAME	SCIENTIFIC NAME
Tigris PS Clethodim	10 to 12 fl. oz./A (0.156-0.188 lb. a.i.)	Tall Fescue	<i>Festuca arundinacea</i>	4 to 6 inches tall (40 to 60% green-up)

**ADJUVANT:** **Tigris PS Clethodim** must be applied with crop oil concentrate at 1 qt./A, plus a spray grade ammonium sulfate at 2.5 to 4 lbs./A.

**MIXING ORDER:** Thoroughly mix spray grade ammonium sulfate in water, add **Tigris PS Clethodim**, then add crop oil concentrate.

#### SPECIAL APPLICATION INSTRUCTIONS / RESTRICTIONS / PRECAUTIONS

Burn or mow fields a minimum of 3 weeks prior to application to remove excess crop residue. Apply in the spring, at 40 to 60% tall fescue green-up, prior to emergence of warm-season grasses. **DO NOT** mow area for 2 weeks after the **Tigris PS Clethodim** application.

Apply in a minimum of 15 to 20 gals. of water per acre at a spray pressure of 40 to 60 PSI at the nozzle.

Apply using flat fan or hollow cone nozzles. **DO NOT** use flood jet nozzles.

Apply only to fields that have warm-season grasses established for 2 years. Applications of **Tigris PS Clethodim** to emerged warm-season grasses may cause injury. **DO NOT** apply to warm-season grasses grown for seed.

**DO NOT** graze treated fields or feed treated forage and/or hay to livestock. **DO NOT** plant any crop for 30 days after application, unless clethodim is registered for use in that crop.

**NOTE:** **Tigris PS Clethodim** applications are most effective if applied when average nighttime temperatures are consistently greater than or equal to 47°F.

Table 20. TIGRIS PS CLETHODIM FOR THE SUPPRESSION OF TALL FESCUE SEED-HEADS IN NON-PRODUCING AGRICULTURAL AREAS

PRODUCT	PRODUCT RATE	SUPPRESSION	APPLICATION TIMING
Tigris PS Clethodim	1.5 to 2 fl. oz./A (0.023-0.031 lb. a.i.)	Tall Fescue Seed-Heads ( <i>Festuca arundinacea</i> )	(50 to 90% Tall Fescue green-up)

**ADJUVANT:** **Tigris PS Clethodim** must be applied with crop oil concentrate at 1 qt./A, plus a spray grade ammonium sulfate at 2.5 to 4 lbs./A.

**MIXING ORDER:** Thoroughly mix spray grade ammonium sulfate in water, add **Tigris PS Clethodim**, then add crop oil concentrate.

#### SPECIAL APPLICATION INSTRUCTIONS / RESTRICTIONS / PRECAUTIONS

Apply at 50 to 90% tall fescue green-up.

Use the higher **Tigris PS Clethodim** rate if less tall fescue green matter is present.

**DO NOT** mow area for 2 weeks after the **Tigris PS Clethodim** application.

Apply in a minimum of 15 to 20 gals. of water per acre at a spray pressure of 40 to 60 PSI at the nozzle. Apply using flat fan or hollow cone nozzles. **DO NOT** use flood nozzles.

2,4-D ester may be added to this tank mix for broadleaf control (see 2,4-D ester label for weeds controlled).

**DO NOT** graze treated fields or feed treated forage and/or hay to livestock. **DO NOT** plant any crop for 30 days after application, unless clethodim is registered for use in that crop.

#### DIRECTIONS FOR USE IN ORNAMENTALS

For ornamental plant uses, **Tigris PS Clethodim** can be used to control labeled grass weeds in greenhouses, bathhouses, shadehouses, and around outdoor ornamentals, including nurseries, parks, roadside plantings, and structure landscapes. The plants listed below have been tested for use with **Tigris PS Clethodim** applications. See tables with listed ornamentals (ornamental trees, ground covers, garden flowers and plants, and shrubs).

Use Rates Per Acre: 6-16 fl. oz. (0.094-0.25 lb. a.i.)

#### Special Use Instructions

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).

Use of crop oil concentrate is not advised since it may injure flowers and foliage.

For repeat applications, make on a minimum of a 14-day interval.

#### IMPORTANT

**Tigris PS Clethodim** successfully controls weeds in newly transplanted and established non-grassy ornamentals. Plant resistance to **Tigris PS Clethodim** at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is advised that the user determine if herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of **Tigris PS Clethodim** have investigated the safety factor to ornamental plants not listed on the label.

The following plants have shown a resistance for **Tigris PS Clethodim** applications:

#### ORNAMENTAL TREES

COMMON NAME	SCIENTIFIC NAME
Alder, Red	<i>Alnus rubra</i>
Ash	<i>Fraxinus</i> spp.
Basswood	<i>Tilia</i> spp.
Birch, European White	<i>Betula pendula</i>
Birch, River	<i>Betula nigra</i>
Birch, White	<i>Betula papyrifera</i>
Crabapple, Flowering	<i>Malus</i> <i>holliana</i>
Dogwood, Flowering	<i>Cornus</i> <i>florida</i>
Golden Chain tree	<i>Laburnum anagyroides</i>
Maples	<i>Acer</i> spp.
Mulberry, White	<i>Morus alba</i>
Oaks	<i>Quercus</i> spp.
Olive, Wild	<i>Olea europaea</i>
Redbud, Eastern	<i>Cercis Canadensis</i>
Sweet Gum, American	<i>Liquidambar styraciflua</i>

#### GROUND COVERS

COMMON NAME	SCIENTIFIC NAME
Bugleweed, Carpet	<i>Ajuga reptans</i>
Ivy, English	<i>Hedera helix</i>
Japanese Spurge	<i>Pachysandra terminalis</i>
Lilyturf	<i>Liriope muscari</i>
Moneywort	<i>Lysimachia nummularia</i>
Mondo Grass, White	<i>Ophiopogon jaburan</i>
Mondo Grass, Dwarf	<i>Ophiopogon japonicus</i>
Periwinkle, Lesser	<i>Vinca minor</i>

GARDEN FLOWERS AND PLANTS

COMMON NAME	SCIENTIFIC NAME
Ageratum	<i>Ageratum</i> spp.
Alyssum*, Sweet	<i>Lobularia maritima</i>
Asparagus Fern	<i>Asparagus setaceus</i>
Bleeding Heart	<i>Dicentra spectabilis</i>
Cast Iron Plant	<i>Aspidistra elatior</i>
Chrysanthemum	<i>Chrysanthemum</i> spp.
Cinquefoil	<i>Potentilla</i> spp.
Coleus	<i>Coleus</i> spp.
Coralbells	<i>Heuchera sanguinea</i>
Cranesbill	<i>Geranium</i> spp.
Dahlia	<i>Dahlia</i> spp.
Daisy, Trailing African	<i>Osteospermum fruticosum</i>
Daylily	<i>Hemerocallis</i> spp.
Dusty Miller	<i>Senecio cineraria</i>
Euonymus	<i>Euonymus</i> spp.
Gazania	<i>Gazania</i> spp.
Geranium, House	<i>Pelargonium hortorum</i>
Heather, False	<i>Cuphea hyssopifolia</i>
Hosta	<i>Hosta fortunei</i>
Iris	<i>Iris</i> spp.
Jasmine Tobacco	<i>Nicotiana alata</i>
Loosestrife	<i>Lythrum salicaria</i>
Margold	<i>Tarotess</i> spp.
Partridgeberry	<i>Mitchella repens</i>
Petunia*	<i>Petunia hybrida</i>
Phlox	<i>Phlox</i> spp.
Pinks	<i>Dianthus</i> spp.
Portulaca	<i>Portulaca grandiflora</i>
Salvia	<i>Salvia</i> spp.
Saxifrage	<i>Saxifrage</i> spp.
Sedum	<i>Sedum</i> spp.
Selloum	<i>Philodendron selloum</i>
Snapdragon*	<i>Antirrhinum majus</i>
Sweet Flag	<i>Acorus gramineus</i>
Tickseed	<i>Coreopsis grandiflora</i>
Touch-Me-Not	<i>Impatiens</i> spp.
Verbena	<i>Verbena</i> spp.
Violet	<i>Viola</i> spp.
Yarrow, Common	<i>Achillea millefolium</i>
Zinnia	<i>Zinnia elegans</i>

\*Slight foliage or flower speckling has been observed on these species.

## SHRUBS

COMMON NAME	SCIENTIFIC NAME
Abelia	<i>Abelia</i> spp.
Anise, Purple	<i>Illicium floridanum</i>
Aucuba	<i>Aucuba</i> spp.
Azalea*	<i>Rhododendron</i> spp.
Bamboo	<i>Bambusa</i> spp.
Barberry, Japanese	<i>Berberis thunbergii</i>
Barberry, Magellan	<i>Berberis buxifolia</i>
Bayberry	<i>Myrica pensylvanica</i>
Bottlebrush	<i>Callistemon citrinus</i>
Boxwood, Common	<i>Buxus sempervirens</i>
Camellia, Common	<i>Camellia japonica</i>
Candytuft	<i>Iberis sempervirens</i>
Cleyera	<i>Cleyera japonica</i>
Coralberry	<i>Ardisia crenata</i>
Crape Myrtle	<i>Lagerstroemia indica</i>
Coyote Brush	<i>Baccharis pilularis</i>
Fig, Creeping	<i>Ficus pumila</i>
Gardenia	<i>Gardenia</i> spp.
Holly	<i>Ilex</i> spp.
Honeysuckle	<i>Lonicera</i> spp.
Indian Hawthorn	<i>Raphiolepis indica</i>
Jasmine	<i>Jasminum</i> spp.
Jasmine, Asiatic	<i>Trachelospermum asiaticum</i>
Jasmine, Star	<i>Trachelospermum jasminoides</i>
Juniper	<i>Juniperus</i> spp.
Lantana	<i>Lantana</i> spp.
Nandina* Bamboo, Heavenly	<i>Nandina domestica</i>
Oleander, Common	<i>Nerium oleander</i>
Oregon Grape	<i>Mahonia aquifolium</i>
Photinia	<i>Photinia</i> spp.
Pittosporum	<i>Pittosporum</i> spp.
Podocarpus	<i>Podocarpus</i> spp.
Privet	<i>Ligustrum</i> spp.
Pyracantha	<i>Pyracantha</i> spp.
Rhododendron	<i>Rhododendron</i> spp.
Rose	<i>Spiraea bumalda</i>
Sweet Olive	<i>Osmanthus fragrans</i>
Viburnum	<i>Viburnum tinus</i>
Wisteria	<i>Wisteria</i> spp.
Yellow Sage/Shrub Verbena	<i>Lantana camara</i>

\*Slight foliage or flower speckling has been observed on these species.

(continued)

**DIRECTIONS FOR ANNUAL GRASSES IN ORNAMENTALS**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass and weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

**Restrictions:**

- **DO NOT** apply more than 16 fl. oz./A (0.25 lb. a.i./A) per application.
- **DO NOT** make more than 4 applications per acre per year.
- **DO NOT** apply more than 64 fl. oz./A (1 lb. a.i./A) per year.
- For repeat applications, make on a minimum of a 14-day interval.

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (Inches)	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE FL. OZ./ACRE <sup>(2)</sup>
Barnyardgrass	<i>Echinochloa crusgalli</i>	2 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Broadleaf Signalgrass	<i>Bracharia platyphylla</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Brome				
California	<i>Bromus carinatus</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Cheat	<i>Bromus secalinus</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Downy	<i>Bromus tectorum</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Ripgut	<i>Bromus diandrus</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Canarygrass	<i>Phalaris canariensis</i>	1 to 4	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Crabgrass				
Hairy	<i>Digitaria adscondens</i>	2 to 6**	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Large	<i>Digitaria sanguinalis</i>	2 to 6**	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Smooth	<i>Digitaria ischaemum</i>	2 to 6**	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Southern	<i>Digitaria ciliaris</i>	2 to 6**	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	2 to 6**	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Fall Panicum	<i>Panicum dichotomiflorum</i>	2 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Field Sandbur	<i>Cenchrus incertus</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Foxtail				
Giant	<i>Setaria faberi</i>	2 to 12	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Green	<i>Setaria viridis</i>	2 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Yellow	<i>Setaria glauca</i>	2 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Goosegrass	<i>Eleusine indica</i>	2 to 6**	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Itchgrass	<i>Rottboellia cochinchinensis</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Junglerice	<i>Echinochloa colona</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)

\* Generally occurs between 3-leaf stage and tillering.

\*\* Length of lateral growth.

(1) 8 fl. oz./A (0.125 lb. a.i./A) = approximately 0.2 fl. oz./1,000 sq. ft.

(2) 16 fl. oz./A (0.25 lb. a.i./A) = approximately 0.4 fl. oz./1,000 sq. ft.

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).

(continued)

**DIRECTIONS FOR ANNUAL GRASSES IN ORNAMENTALS (cont'd)**

GRASS SPECIES	SCIENTIFIC NAME	WEED HEIGHT* (Inches)	RATE FL. OZ./ACRE(1)	HIGH RATE FL. OZ./ACRE(2)
Lovegrass (Stinkgrass)	<i>Eragrostis cilianensis</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Rabbitsfootgrass	<i>Polypogon monspeliensis</i>	1 to 4	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Red Rice	<i>Oryza sativa</i>	1 to 3	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Ryegrass				
Hardy	<i>Lolium remontum</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Italian	<i>Lolium multiflorum</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Seedling Johnsongrass	<i>Sorghum halepense</i>	4 to 10	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Shattercane	<i>Sorghum bicolor</i>	6 to 18	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Southwestern Cupgrass	<i>Eriochloa gracilis</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Spangletop				
Amazon	<i>Leptochloa panicoides</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Bearded	<i>Leptochloa fascicularis</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Mexican	<i>Leptochloa uninervia</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Red	<i>Leptochloa filiformis</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Texas Panicum	<i>Panicum texanum</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Volunteer Cereals				
Barley	<i>Hordeum vulgare</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Oats	<i>Avena sativa</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Rye	<i>Secale cereale</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Wheat	<i>Triticum aestivum</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Volunteer Corn	<i>Zea mays</i>	4 to 12	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Volunteer Corn	<i>Zea mays</i>	12 to 24	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Volunteer Grain Sorghum	<i>Sorghum bicolor</i>	8 to 12	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Wild Oats	<i>Avena fatua</i>	2 to 6	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Wild Proso Millet	<i>Panicum miliaceum</i>	2 to 10	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Witchgrass	<i>Panicum capillare</i>	2 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Woolly Cupgrass	<i>Eriochloa villosa</i>	2 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)

\* Generally occurs between 3-leaf stage and tillering.

\*\* Length of lateral growth.

(1) 8 fl. oz./A (0.125 lb. a.i./A) = approximately 0.2 fl. oz./1,000 sq. ft.

(2) 16 fl. oz./A (0.25 lb. a.i./A) = approximately 0.4 fl. oz./1,000 sq. ft.

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gal. (0.25% v/v).

**DIRECTIONS FOR ANNUAL BLUEGRASS CONTROL WITH TIGRIS PS CLETHODIM IN ORNAMENTALS**

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE FL. OZ./ACRE
Annual Bluegrass ( <i>Poa annua</i> )	to 4-leaf	6 (0.094 lb. a.i.)	16 (0.25 lb. a.i.)

Apply under favorable soil moisture and humidity that exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).

Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.

Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).

**Restrictions:**

- **DO NOT** apply more than 16 fl. oz./a (0.25 lb. a.i./a) per application.
- **DO NOT** make more than 4 applications at the lowest rate, 6 fl. oz. (0.094 lb. a.i./a), or 2 applications at the highest rate, 16 fl. oz. (0.25 lb. a.i./a), per acre per year.
- For repeat applications, make on a minimum of a 14-day interval.
- **DO NOT** apply more than 32 fl. oz./a (0.5 lb. a.i./a) per year.

**DIRECTIONS FOR PERENNIAL GRASSES**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

**Restrictions:**

- **DO NOT** apply more than 16 fl. oz./a (0.25 lb. a.i./a) per application.
- **DO NOT** make more than 2 applications per acre per year.
- For repeat applications, make on a minimum of a 14-day interval.
- **DO NOT** apply more than 32 fl. oz./a (0.5 lb. a.i./a) per year.

GRASS SPECIES	WEED HEIGHT (inches)	RATE FL. OZ./ACRE <sup>(1)</sup>	HIGH RATE FL. OZ./ACRE (2)
Bermudagrass ( <i>Cynodon dactylon</i> )			
First Application	3 (or up to 6" runners)	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	3 (or up to 6" runners)	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Quackgrass ( <i>Elytrigia repens</i> )			
First Application	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Rhizome Johnsonsgrass ( <i>Sorghum halepense</i> )			
First Application	12 to 24	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	6 to 18	6 (0.094 lb. a.i.)	8 (0.125 lb. a.i.)
Wirestem Muhy ( <i>Muhlenbergia frondosa</i> )			
First Application	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)
Repeat Application(s) (if regrowth occurs)	4 to 8	8 (0.125 lb. a.i.)	16 (0.25 lb. a.i.)

(1) 8 fl. oz./a (0.125 lb. a.i./a) = approximately 0.2 fl. oz./1,000 sq. ft.

(2) 16 fl. oz./a (0.25 lb. a.i./a) = approximately 0.4 fl. oz./1,000 sq. ft.

Add a non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

### PESTICIDE STORAGE:

Keep pesticide in original container. **DO NOT** put concentrate or dilute into food or drink containers. Store in cool, dry place. **DO NOT** store diluted spray.

### PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**[Nonrefillable plastic container < 5 gallons:] Nonrefillable container.** **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinse into application equipment or a mix tank or store rinse for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**[Nonrefillable plastic container > 5 gallons:] Nonrefillable container.** **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinse into application equipment or a mix tank or store rinse for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**[Nonrefillable metal container > 5 gallons:] Nonrefillable container.** **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinse into application equipment or a mix tank or store rinse for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

**[Refillable Containers:] Refillable container.** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinse into application equipment or rinse collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Tigris, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Tigris, LLC and Seller harmless for any claims relating to such factors.

Tigris, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Tigris, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, TIGRIS, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Tigris, LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF TIGRIS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF TIGRIS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Tigris, and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Tigris, LLC.

All trademarks are the property of their respective owners.



# TIGRIS™ PS CLETHODIM

## HERBICIDE

CLETHODIM GROUP 1 HERBICIDE

**ACTIVE INGREDIENT:**

\*Clethodim..... 26.4%

**OTHER INGREDIENTS:**

**TOTAL:**..... 73.6%

Contains Petroleum Distillates

\* $(E)$ -2-[[( $(S$ )-chloro-2-propenyl)-oxy]mino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one

Contains 2.0 lbs. clethodim per gallon.

### KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende lo etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>Immediately call a poison control center or doctor.</li> <li><b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li><b>DO NOT</b> give any liquid to the person.</li> <li><b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

#### HOTLINE NUMBERS

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at **1-800-858-7378**, Mon.-Fri. 8:00 a.m. to 12:00 p.m. Pacific Time or your poison control center at **1-800-222-1222**.

#### NOTE TO PHYSICIAN

Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid, which can cause pneumonitis. If ingested, probable mucosal damage may contraindicate the use of gastric lavage.

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

##### WARNING

Causes substantial but temporary eye irritation. Avoid contact with skin. **DO NOT** get in eyes, on skin, or on clothing. Harmful if swallowed or inhaled. Avoid breathing vapors or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### ENVIRONMENTAL HAZARDS

**DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply where runoff is likely to occur. **DO NOT** apply where weather conditions favor drift from areas treated. **DO NOT** contaminate water when disposing of equipment wash water or rinses.

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice. Use of this product is prohibited in the following areas where the species are known to exist.

• **Solano Grass:** Solano County, California: the vernal lakes are bounded by the Union Pacific Railroad and Hastings Road to the north, Highway 113 to the east, Highway 12 to the south, and Travis Air Force Base to the west.

• **Wild Rice:** Yolo County, Texas.

#### PHYSICAL OR CHEMICAL HAZARDS

**Combustible.** **DO NOT** use or store near heat or open flame.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

#### PESTICIDE STORAGE:

Keep pesticide in original container. **DO NOT** put concentrate or dilute into food or drink containers. Store in cool, dry place. **DO NOT** store diluted spray.

#### PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER HANDLING:

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

See label booklet for complete Precautionary Statements and Directions for Use.

EPA Reg. No.: 92647-31

EPA Est. No.: 70989-MO-001 (OS)

05905-IA-001 (HD)

Letter(s) in the lot number correspond to letter(s) following the EPA Est. No.



#### Manufactured For:

Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557